arXiv:1204.2176v1 [math.PR] 10 Apr 2012

LARGE TIME ASYMPTOTICS FOR THE FLUCTUATION SPDE IN THE KURAMOTO SYNCHRONIZATION MODEL

ERIC LUÇON[1]

Abstract. We investigate the long-time asymptotics of the fluctuation SPDE in the Kuramoto synchronization model. We mainly establish the linear behavior for large time and weak disorder of the quenched limit fluctuations of the empirical measure of the oscillators around its McKean-Vlasov limit. This is carried out through a precise spectral analysis of the unbounded evolution operator, using arguments of perturbations of self-adjoint operators and analytic semigroups. We state in particular a Jordan decomposition of the evolution operator which is the key point in order to show that the fluctuations of the disordered Kuramoto model are not self-averaging.

1. Introduction

1.1. **Synchronization of heterogeneous oscillators.** Collective behavior of oscillators and synchronization phenomenon are the subject of a vast literature either in biological (neuronal models, collective behavior of insects, cells, etc.) or in physical contexts (see [25, 24, 40] and references therein). While a precise description of each of the different instances in which synchronization emerges demands specific, possibly very complex models, the celebrated Kuramoto model [1, 39] has emerged as capturing some of the fundamental aspects of synchronization.

The *disordered Kuramoto* model concerns a family of heterogeneous oscillators (or rotators) on the circle $\mathbf{S} := \mathbf{R}/2\pi\mathbf{Z}$ in a noisy mean-field interaction (that is the interaction is perturbed by a thermal noise), in the presence of disorder. More precisely, we make the assumption that the rotators are similar but not necessarily identical. In that sense, the model captures the fact that each rotator has a tendency to obey to its own natural frequency which may differ from one rotator to another. Those frequencies are chosen at random and independently for each rotator according to a probability distribution $\mu(\,\mathrm{d}\omega)$ on $\mathbf{R}$; hence, this supplementary source of randomness will be considered as a *disorder*. One of the main characteristics of the Kuramoto model is that it presents a phase transition, as the coupling strength between the rotators increases, from an incoherent state where the rotators are not synchronized to a synchronous one where the phases of the rotators concentrate around a common value.

1.1.1. *The continuous model.* The disordered Kuramoto model [25, 1], in the limit of an infinite population of rotators, is given by the following system of coupled nonlinear partial

---

[1] Université Pierre et Marie Curie (Paris 6) and Laboratoire de Probabilités et Modèles Aléatoires (CNRS), U.F.R. Mathématiques, Case 188, 4 place Jussieu, 75252 Paris cedex 05, France

Date: March 2, 2013.

2010 *Mathematics Subject Classification.* 35P15, 46N60, 47A55, 60H15, 82C22.

Key words and phrases. Stochastic partial differential equations - perturbation of analytic operators - Jordan decomposition - Kuramoto model - synchronization - disordered systems - self-averaging.

differential equations (*McKean-Vlasov equations*): for all ω in $\mathrm{Supp}(\mu)$

$$\partial_t q_t(\theta,\omega) = \frac{1}{2}\partial_\theta^2 q_t(\theta,\omega) - \partial_\theta\Big(q_t(\theta,\omega)\left(\langle J * q_t\rangle_\mu(\theta) + \omega\right)\Big), \quad t>0,\ \theta\in\mathbf{S}, \tag{1.1}$$

with periodic boundary conditions and initial condition given by

$$\forall\omega\in\mathrm{Supp}(\mu), \quad q_t(\theta,\omega)\,\mathrm{d}\theta \xrightarrow[t\searrow 0]{} \gamma(\,\mathrm{d}\theta) \tag{1.2}$$

for some probability law γ on the circle $\mathbf{S}$. Here,

$$\langle J * q_t\rangle_\mu(\theta) = -K\int_{\mathbf{R}}\int_{\mathbf{S}} \sin(\varphi) q_t(\theta-\varphi,\omega)\,\mathrm{d}\varphi\mu(\,\mathrm{d}\omega), \tag{1.3}$$

stands for the convolution of $J(\cdot) := -K\sin(\cdot)$ with q_t, averaged with respect to the disorder.

Note that we are looking for solutions $(t,\theta,\omega)\mapsto q_t(\theta,\omega)$ to (1.1)–(1.2) such that $q_t(\cdot,\omega)$ is a probability density for all fixed t and ω: $q_t(\cdot,\cdot)\geqslant 0$ for all $t>0$ with $\int_{\mathbf{S}} q_t(\theta,\omega)\,\mathrm{d}\theta = 1$ for all $\omega\in\mathrm{Supp}(\mu)$. In that sense, the physical interpretation of (1.1)–(1.2) is the following: $\theta\in\mathbf{S}\equiv[0,2\pi]$ is the phase of the rotators, $\gamma(\,\mathrm{d}\theta)$ is the initial distribution of the rotators on $\mathbf{S}$, $K>0$ is a nonnegative constant representing the coupling strength between the rotators, μ is the probability distribution of the frequencies, and $q_t(\theta,\omega)$ is the density at time $t>0$ of rotators with phase θ and frequency ω.

We stress the fact that evolution (1.1) exhibits the following properties: it has quadratic nonlinearity, it is nonlocal, and if the support of μ is infinite, (1.1) is in fact equivalent to an infinite number of coupled equations; indeed, in order to know $q(\cdot,\omega)$ for one $\omega\in\mathrm{Supp}(\mu)$, it is necessary to know $q(\cdot,\omega')$ for every $\omega'\in\mathrm{Supp}(\mu)$. To fix ideas, one may for example consider the simple binary case where $\mu=\frac{1}{2}(\delta_{-1}+\delta_1)$. Then (1.1) reduces to two equations (one for $+1$, the other for -1) which are coupled via the averaged probability measure $\frac{1}{2}(q_t(\theta,+1)+q_t(\theta,-1))\,\mathrm{d}\theta$. But for more general situations ($\mu=\mathcal{N}(0,1)$ say) this would consist of an infinite number of coupled equations.

Remark 1.1. *It is immediate to see that* (1.1) *exhibits the following symmetries:*

- Rotation invariance: *if* $q_t(\theta,\omega)$ *solves* (1.1) *so does* $q_t(\cdot+\theta_0,\omega)$ *for any constant* $\theta_0\in\mathbf{S}$*,*
- Even symmetry: *if* $q_t(-\theta,-\omega)|_{t=0} = q_t(\theta,\omega)|_{t=0}$*, then it is true for all* $t>0$*.*

In particular, the stationary solutions of (1.1) *will share these symmetries, see* (1.13)*.*

The Kuramoto model (1.1), as well as many generalizations, has been widely discussed mostly in the physical literature. We refer to [1, 39] and references therein for a global review on the subject. We highlight in particular the *active rotators model* (see e.g. [37]) which concerns similar interacting rotators in the presence of an additional potential. The dynamical properties for this model tend to be fairly more complicated than the model studied here. We refer to a recent result [19] that proves in particular the existence and stability of invariant normally hyperbolic manifolds for the active rotators model.

Note that uniqueness of a solution to (1.1) follows from standard arguments concerning fundamental solutions of parabolic equations ([2, 15]) and has been rigorously established in [17, § A]. Another proof of uniqueness can be found in [10] on the basis of heat kernel estimates under regularity assumptions on the initial condition.

Existence of a solution to (1.1) may be seen as a consequence of the following probabilistic interpretation of (1.1).

1.1.2. *The microscopic model.* The system (1.1) has a natural microscopic interpretation which is the following: for all $N \geqslant 1$, pick up independently at random a sequence of frequencies $\omega_1, \dots, \omega_N \in \mathbf{R}$ according to the distribution μ and consider the following system of N stochastic differential equations (whose solutions $(\theta_j)_{1 \leqslant j \leqslant N}$ are identified with their projection on $\mathbf{S}$ $(\theta_j \mod 2\pi)_{1 \leqslant j \leqslant N}$):

$$\forall j = 1, \dots, N, \ \mathrm{d}\theta_{j,t} = \omega_j \,\mathrm{d}t - \frac{K}{N} \sum_{i=1}^{N} \sin(\theta_{j,t} - \theta_{i,t}) \,\mathrm{d}t + \mathrm{d}B_{j,t}, \tag{1.4}$$

where at time $t = 0$, the rotators θ_j are i.i.d. samples of the law γ and $\{B_j\}_{j=1,\dots,N}$ is a sequence of N standard independent Brownian motions (which stands for the thermal noise in the system). We stress the fact that we have here *two sources of randomness*: the randomness that chooses the frequencies $(\omega_j)_{j \geqslant 1}$ and the randomness coming from the Brownian motions. Hence, one can see (1.4) as a particular instance of interacting diffusions in a random environment.

Evolution (1.1) appears naturally as the large N-limit of the system (1.4) in the following way: let us define the empirical probability measure ν_N of both rotators and frequencies:

$$t \mapsto \nu_{N,t} := \frac{1}{N} \sum_{j=1}^{N} \delta_{(\theta_{j,t}, \omega_j)} \in \mathcal{C}([0, +\infty), \mathcal{M}_1(\mathbf{S} \times \mathbf{R})), \tag{1.5}$$

where $\delta_{(\theta,\omega)}$ is the Dirac measure in (θ, ω) and $\mathcal{M}_1(\mathbf{S} \times \mathbf{R})$ the set of probability measures on $\mathbf{S} \times \mathbf{R}$. It can be shown (see [10, 26]) that, under mild hypotheses, the sequence ν_N converges in law (as a process), as N goes to $+\infty$, to the deterministic limit $t \mapsto \nu_t$ such that

$$\begin{cases} \nu_0(\mathrm{d}\theta, \mathrm{d}\omega) &=& \gamma(\mathrm{d}\theta) \otimes \mu(\mathrm{d}\omega) \\ \nu_t(\mathrm{d}\theta, \mathrm{d}\omega) &=& q_t(\theta, \omega) \,\mathrm{d}\theta \mu(\mathrm{d}\omega), \quad t > 0, \end{cases} \tag{1.6}$$

where q_t is solution of the McKean-Vlasov equation (1.1).

Remark 1.2. *We stress the fact that the above convergence holds* for almost every choice of the initial sequence of frequencies $(\omega_j)_{j \geqslant 1}$ *(see* [26, Th. 2.5]*); due to the mean-field character of the diffusion* (1.4)*, there is a* self-averaging phenomenon*: the limit as $N \to +\infty$ given by* (1.1) *is deterministic and does not depend on the initial choice of the disorder $(\omega_j)_{j \geqslant 1}$.*

The above convergence is a disordered generalization of known results about mean-field interacting diffusions. One can cite e.g. [16, 21, 22, 29, 32] for similar problematics concerning models without disorder.

1.2. **The fluctuation SPDE.** In this paper, we investigate the asymptotic behavior as $t \to +\infty$ of the following stochastic partial differential equation:

$$\eta_t = \eta_0 + \int_0^t L_{q_s} \eta_s \,\mathrm{d}s + W_t, \tag{1.7}$$

where L_{q_s} is the linear operator arising from the linearization around the solution $t \mapsto q_t$ of nonlinear evolution (1.1):

$$L_{q_t} h(\theta, \omega) := \frac{1}{2} \partial_\theta^2 h(\theta, \omega) - \partial_\theta \left(h(\theta, \omega) \left(\langle J * q_t \rangle_\mu (\theta) + \omega \right) + q_t(\theta, \omega) \langle J * h \rangle_\mu (\theta) \right), \tag{1.8}$$

and where W is a Gaussian process, indexed by functions $\varphi(\theta,\omega)$ such that $\partial_\theta\varphi(\cdot,\omega)\in\mathbf{L}^2(\mathbf{S})$ for all $\omega\in\mathrm{Supp}(\mu)$, with covariance

$$\forall\varphi_1,\varphi_2,\ \mathbf{E}(W_t(\varphi_1)W_s(\varphi_2))=\int_0^{s\wedge t}\int_{\mathbf{S}\times\mathbf{R}}\partial_\theta\varphi_1(\theta,\omega)\partial_\theta\varphi_2(\theta,\omega)q_u(\theta,\omega)\,\mathrm{d}\theta\mu(\,\mathrm{d}\omega)\,\mathrm{d}u.\tag{1.9}$$

As we will see later in the introduction, the SPDE (1.7) is the natural limit object in the Central Limit Theorem associated to the convergence of the empirical measure ν_N (1.5) towards its limit ν (1.6), as $N\to+\infty$. Namely, the object of a previous work [26] was to prove that the fluctuation process

$$\eta_N:=\sqrt{N}\,(\nu_N-\nu)\tag{1.10}$$

converges, in a weak sense, in an appropriate space of distributions on $\mathbf{S}\times\mathbf{R}$, to the solution η of (1.7). The main motivation for this work was to study the influence of a typical choice of the disorder $(\omega_j)_{j\geqslant 1}$ (*quenched model*) on the finite-size behavior of the microscopic Kuramoto model (1.4). As a matter of fact, it turns out that the limit process η illustrates the dependence of the system (1.4) on the fluctuations of the frequencies $(\omega_j)_{j\geqslant 1}$: as we will see, fluctuations of the disorder in (1.4) compete with the fluctuations of the thermal noise and make the whole system rotate. What makes evolution (1.7) relevant here is that its solution η still captures this disorder-dependent rotation; at least numerically, one observes trajectories of the process η that are compatible with the ones observed for the finite-size system (1.4) (see Figures 2 and 3).

Hence, the main result of the paper (Theorem 2.9) concerns the issue of the asymptotics of the solution η of (1.7) as time t goes to infinity. More precisely, the purpose of Theorem 2.9 is to prove the following convergence, in an appropriate space of distribution, as $t\to+\infty$:

$$\frac{\eta_t}{t}\xrightarrow[t\to+\infty]{\text{in law}}v,\tag{1.11}$$

where the law of the limit v (that depends randomly on the disorder) is nontrivial and will be made explicit. In particular, this convergence might be an illustration of the fact that the self-averaging phenomenon for the law of large numbers observed in § 1.1.2 no longer holds at the scale of the central limit theorem: fluctuations in (1.4) depend on the fluctuations of the disorder. Nevertheless, a rigorous derivation of the dependence in the disorder of the microscopic system (1.4) would require to study this system for larger time scales (that is for time scales of order $\sqrt{N}$ instead of time scales of order 1, as it is done in this work). More precisely, one would need to study the *joint* limit of the empirical measure ν_N, as t *and* N goes to infinity. This has not been carried out here[1] and would be a natural perspective for this work.

The result we present here relies on a detailed spectral analysis of the unbounded evolution operator L_q defined in (1.8), using arguments from perturbation theory of self-adjoint operators and of analytic semigroups. The main ingredient for this result is a precise understanding of the structure of the eigenspace of L_q associated to the eigenvalue 0, proving in particular the existence of a nontrivial Jordan block for the eigenvalue 0, relying on *a priori* estimates on the Dirichlet form associated to L_q and an extension of Lax-Milgram Theorem.

Before going into the details of the asymptotics of (1.7), let us motivate our result in view of the recent advances made on the Kuramoto model.

---

[1] In this work, we first make $N\to\infty$ and *then* make $t\to\infty$.

1.3. **Motivations and problems.**

1.3.1. *Notion of synchronization in the Kuramoto model.* It is now well understood (see [36]) that crucial features of evolution (1.1) are captured by the order parameters $r_t \geqslant 0$ and $\psi_t \in \mathbf{S}$ defined by:

$$r_t e^{i\psi_t} = \int_{\mathbf{S}\times\mathbf{R}} e^{i\theta} q_t(\theta,\omega)\,\mathrm{d}\theta\,\mathrm{d}\mu(\omega). \tag{1.12}$$

The quantity r_t captures in fact the degree of synchronization of a solution (the profile $q_t \equiv \frac{1}{2\pi}$ for example corresponds to $r_t = 0$ and represents a total lack of synchronization) and ψ_t identifies the center of synchronization: this is true and rather intuitive for unimodal profiles. In this framework, it turns out that synchronization reads in the existence of nontrivial stationary solutions q to (1.1), which can be parameterized by the order parameter r defined in (1.12): namely, following [36] (see also [11, p. 117]), if μ is symmetric, any equilibrium in (1.1) may be written as $q(\theta+\theta_0,\omega)$ for any fixed arbitrary $\theta_0 \in \mathbf{S}$ where

$$q(\theta,\omega) := \frac{S(\theta,\omega,2Kr)}{Z(\omega,2Kr)}, \tag{1.13}$$

for

$$S(\theta,\omega,x) := e^{G(\theta,\omega,x)}\left[(1-e^{4\pi\omega})\int_0^{\theta} e^{-G(u,\omega,x)}\,\mathrm{d}u + e^{4\pi\omega}\int_0^{2\pi} e^{-G(u,\omega,x)}\,\mathrm{d}u\right], \tag{1.14}$$

where $G(u,\omega,x) = x\cos(u) + 2\omega u$, $Z(\omega,x) = \int_{\mathbf{S}} S(\theta,\omega,x)\,\mathrm{d}\theta$ a normalization constant in order to insure that q is indeed a probability density on $\mathbf{S}$, and $r \in [0,1]$ satisfies (1.12) with q given by (1.13):

$$r = \Psi_\mu(2Kr), \quad \text{where} \quad \Psi_\mu(x) := \int_{\mathbf{R}} \frac{\int_{\mathbf{S}}\cos(\theta) S(\theta,\omega,x)\,\mathrm{d}\theta}{Z(\omega,x)}\mu(\,\mathrm{d}\omega). \tag{1.15}$$

Discriminating between the possible solutions $r \in [0,1]$ to the fixed-point relation (1.15), one has two possibilities:

- The case $r = 0$ is always a solution to (1.15) and corresponds to the constant density $q \equiv \frac{1}{2\pi}$,
- Any solution q with $r > 0$ is called a synchronized (nontrivial) solution. Any easy calculation of the derivative of $\Psi_\mu(\cdot)$ at 0 shows that such solutions exist at least when the coupling strength K is greater than $\tilde{K} := \left(\int_{\mathbf{R}} \frac{\mu(\,\mathrm{d}\omega)}{1+4\omega^2}\right)^{-1}$. In that case, due to the rotation invariance (Remark 1.1), each solution $r > 0$ of (1.15) corresponds to a whole circle of synchronized stationary solutions

$$M_{sync} := \{q(\cdot+\theta_0,\omega);\, \theta_0 \in \mathbf{S}\}. \tag{1.16}$$

1.3.2. *The case with no disorder.* In order to motivate our results, let us give a brief review of the case where there is no disorder, that is $\mu = \delta_0$. In that case, (1.1) reduces to:

$$\partial_t q_t(\theta) = \frac{1}{2}\partial_\theta^2 q_t(\theta) - \partial_\theta\left(q_t(\theta)(J * q_t)(\theta)\right). \tag{1.17}$$

As well as in the disordered case, any non-trivial stationary profile can be written as $q_0(\theta+\theta_0)$ where

$$q_0(\theta) := \frac{e^{2Kr_0\cos(\theta)}}{\int_{\mathbf{S}} e^{2Kr_0\cos(u)}\,\mathrm{d}u} = \frac{e^{2Kr_0\cos(\theta)}}{Z_0(2Kr_0)}, \tag{1.18}$$

where ($K > 1$) r_0 is the only nonnegative solution of

$$r_0 = \Psi_0(2Kr_0), \text{ where } \Psi_0(x) := \frac{\int_{\mathbf{S}} \cos(\theta) e^{x\cos(\theta)}\,\mathrm{d}\theta}{\int_{\mathbf{S}} e^{x\cos(\theta)}\,\mathrm{d}\theta}. \tag{1.19}$$

In that case, since Ψ_0 is strictly concave ([34, Lem. 4]) and $\partial_{r_0}\Psi_0(2Kr_0) = K$, the phase transition is obvious: for $K \leqslant 1$, $r_0 = 0$ is the only solution to (1.19) and $\frac{1}{2\pi}$ is the only stationary solution whereas for $K > 1$ this solution coexists with a unique (up to rotation) synchronized solution (corresponding to the unique $r_0 > 0$ solution to (1.19)).

Recent advances have been made concerning the dynamical properties of the non-disordered Kuramoto model. In particular in [5], the linear stability of the synchronized solution (1.18) is proved, with explicit spectral gap estimates. We will rely on these results for the rest of this paper and we refer to § 2.3 for more details. What is more, the whole long-time dynamics of (1.17) is addressed in [18], discussing in particular the existence and the nature of a global attractor in both cases $K \leqslant 1$ and $K > 1$ and proving non-linear stability of synchronized solutions for $K > 1$.

1.3.3. *Perturbing the non disordered case.* A natural issue is then to understand in what way adding the disorder $(\omega_j)_{j\geqslant 1}$ may perturb the results mentioned for the non-disordered case. In this direction, the question of linear and nonlinear stability of synchronized solutions in the disordered case is addressed in [17], at least in the case where the disorder is small. In that extent, one can distinguish two regimes:

- The case μ not symmetric: from an intuitive point of view, in this case, there are more rotators that are likely to rotate in one given direction than in the other direction; the whole system has a tendency to rotate in this direction. This phenomenon can be made rigorous and reads in fact on the continuous model (1.1): for small disorder, the existence of stable rotating-waves which solve (1.1) is proved in [17], with explicit computation of the speed of rotation.
- The case μ symmetric: in this case, the speed of rotation of the rotating waves in the continuous model (1.1) is equally zero. Nevertheless, one can prove ([17]) by perturbations arguments the linear and nonlinear stability of synchronized stationary solutions, once again for small disorder (with explicit estimates on the size of the disorder).

1.3.4. *Finite system: disorder-induced rotation and non self-averaging phenomenon.* As already mentioned, the study of the SPDE (1.7) is initially motivated by the Central Limit Theorem associated to the convergence of the empirical measure ν_N defined in (1.5) and is linked with the behavior of the microscopic system (1.4) for large but finite N. Indeed, Balmforth and Sassi [4] pointed out on the basis of numerical computations the following remarkable phenomenon: even if the distribution μ is symmetric, the fluctuations of a fixed chosen sample of the disorder $(\omega_1, \ldots, \omega_N)$ makes it *not symmetric* and induces a slow rotation of the center of the synchronization in one direction at a speed of order $N^{-1/2}$ that depends randomly on the sample of the disorder (Fig. 1 and 2). To fix ideas, one may think of the simple case where the distribution of the disorder is binary: $\mu = \frac{1}{2}(\delta_{-1} + \delta_1)$. In a finite sample $(\omega_1, \ldots, \omega_N)$ of spins in $\{\pm 1\}^N$, there may be a majority of positive spins (equal to $+1$) with respect to the negative spins (equal to -1), so that the rotators with positive frequency have a tendency to induce a global rotation in their sense. This non self-averaging phenomenon at the scale of fluctuations (to compare with Remark 1.2) can be noticed by



Figure 1. We plot here the evolution of the marginal on $\mathbf{S}$ of ν_N for $N=600$ oscillators ($\mu=\frac{1}{2}(\delta_{-1}+\delta_1)$, $K=6$). The oscillators are initially chosen independently and uniformly on $[0,2\pi]$ independently of the disorder. First the dynamics leads to synchronization of the oscillators ($t=6$) to a profile which is close to a nontrivial stationary solution of McKean-Vlasov equation (1.1). Secondly, we observe that the center $\psi_{N,t}$ of this density moves to the right with an approximately constant speed; what is more, this speed of fluctuation turns out to be sample-dependent (see Fig. 2).

computing the finite-size order parameters $(r_{N,t},\psi_{N,t})$ (that is the microscopic equivalents of the macroscopic order parameters (r_t,ψ_t) defined in (1.12)):

$$r_{N,t}e^{i\psi_{N,t}}=\frac{1}{N}\sum_{j=1}^{N}e^{i\theta_{j,t}}=\int_{\mathbf{S}\times\mathbf{R}}e^{i\theta}\,\mathrm{d}\nu_{N,t}(\theta,\omega), \tag{1.20}$$

where ν_N is the empirical measure of the system defined in (1.5). In particular, the order parameter $\psi_{N,t}$ captures the position of the center of synchronization for finite N (see Figure 1). It is seen on Figure 2 that $t\mapsto\psi_{N,t}$ has an approximately linear behavior whose slope depends on the choice of the disorder.

1.3.5. *Central Limit Theorem.* The first step to tackle this phenomenon was to prove (see [26]) that the following convergence holds

$$\eta_N \xrightarrow[N\to\infty]{} \eta \tag{1.21}$$

in the space of continuous processes with values in $\mathcal{S}'(\mathbf{S}\times\mathbf{R})$ where η is the solution to (1.7)

$$\eta_t=\eta_0+\int_0^t L_{q_s}\eta_s\,\mathrm{d}s+W_t,$$

and where

- $\mathcal{S}'(\mathbf{S}\times\mathbf{R})$ is understood as the usual space of tempered distributions on $\mathbf{S}\times\mathbf{R}$,
- W is the Gaussian process whose covariance is given in (1.9),
- L_{q_t} is the operator defined in (1.8),



FIGURE 2. Trajectories of the center of synchronization ψ_N for different realizations of the disorder ($\mu = \frac{1}{2}(\delta_{-0.5} + \delta_{0.5})$, $K = 4$, $N = 400$). We observe here the non self-averaging phenomenon: direction and speed of the center depend on the choice of the initial N-sample of the disorder. Moreover, these simulations are compatible with speeds of order $N^{-1/2}$.

- the initial condition η_0 is independent with W.

It is important to note that the convergence that is considered in (1.21) is more elaborate than a usual convergence in law, since this convergence is understood in a quenched set-up, that is for fixed choice of frequencies $(\omega_j)_{j\geqslant 1}$. In fact, the precise notion of convergence used in [26] for (1.21) is not really relevant for the purpose of this paper. The interested reader may find more details in [26, Th. 2.10]. What we only need to retain here is that, in view of the fact that there are two sources of randomness in the microscopic model (1.4), the process η also depends on two sources of randomness at the limit. More precisely, the limit object η captures the dependence of the disorder through its nontrivial mean-value: there exists a Gaussian process $\omega \mapsto C(\omega)$ with covariance

$$\forall \varphi_1, \varphi_2 : \mathbf{S}\times\mathbf{R} \to \mathbf{R},\ \Gamma_C(\varphi_1, \varphi_2) = \mathrm{Cov}_\mu\left(\int_{\mathbf{S}} \varphi_1(\cdot, \omega)\,\mathrm{d}\gamma, \int_{\mathbf{S}} \varphi_2(\cdot, \omega)\,\mathrm{d}\gamma\right) \tag{1.22}$$

such that for fixed ω, the initial condition $\eta_0 = \eta_0^\omega$ in (1.7) may be written as

$$\eta_0^\omega = X + C(\omega), \tag{1.23}$$

where X is a centered Gaussian process with explicit covariance, and $C(\omega)$ is the constant mean-value of η_0^ω. Let us give a microscopic intuition of the decomposition (1.23): at time

$t = 0$, for $N \geqslant 1$ and $\varphi : \mathbf{S} \times \mathbf{R} \to \mathbf{R}$, $\eta_{N,0}(\varphi)$ defined in (1.10) may be decomposed into

$$\begin{aligned}\eta_{N,0}(\varphi) &= \frac{1}{\sqrt{N}} \sum_{j=1}^{N} \Big(\varphi(\theta_j, \omega_j) - \int_{\mathbf{S}\times\mathbf{R}} \varphi(\theta, \omega) \gamma(\mathrm{d}\theta) \mu(\mathrm{d}\omega) \Big), \\ &= \frac{1}{\sqrt{N}} \sum_{j=1}^{N} \Big(\varphi(\theta_j, \omega_j) - \int_{\mathbf{S}} \varphi(\theta, \omega_j) \gamma(\mathrm{d}\theta) \Big) \\ &+ \frac{1}{\sqrt{N}} \sum_{j=1}^{N} \Big(\int_{\mathbf{S}} \varphi(\theta, \omega_j) \gamma(\mathrm{d}\theta) - \int_{\mathbf{S}\times\mathbf{R}} \varphi(\theta, \omega) \gamma(\mathrm{d}\theta) \mu(\mathrm{d}\omega) \Big), \\ &:= X_N(\varphi) + C_N(\varphi).\end{aligned} \tag{1.24}$$

In particular, by definition of X_N, this process is centered for all choice of $(\omega_j)_{j\geqslant 1}$ and converges to some X for almost every $(\omega_j)_{j\geqslant 1}$ ([6, Th. 27.3]). As far as the process C_N is concerned, it is easily seen that it converges in law (w.r.t. the disorder) to the process C with covariance given by (1.22). What is important to understand here is that the random variable $C(\cdot)$ precisely captures the fluctuations of the disorder. For example, let us consider the case where $\mu = \frac{1}{2}(\delta_{-\omega_0} + \delta_{\omega_0})$, for some $\omega_0 > 0$. As we will see in Theorem 2.9, the relevant quantity is in fact the process C_+ (indexed by functions $\psi : \mathbf{S} \to \mathbf{R}$) that is the restriction of the process C to the component on $+\omega_0$:

$$\forall \psi : \mathbf{S} \to \mathbf{R},\ C_{+,\psi} := C_{\psi \mathbb{1}_{\omega=\omega_0}}. \tag{1.25}$$

Thanks to (1.24), C_+ is the limit in law of the microscopic process $C_{N,+}$ (indexed by functions $\psi : \mathbf{S} \to \mathbf{R}$) defined by

$$\forall \psi,\ C_{N,+}(\psi) := \Big(\int_{\mathbf{S}} \psi(\cdot)\, \mathrm{d}\gamma \Big) \frac{1}{\sqrt{N}} \sum_{i=1}^{N} \Big(\mathbb{1}_{(\omega_i=\omega_0)} - \frac{1}{2} \Big) := \Big(\int_{\mathbf{S}} \psi(\cdot)\, \mathrm{d}\gamma \Big) \frac{\alpha_N}{\sqrt{N}}. \tag{1.26}$$

Here, α_N is exactly the (centered) number of frequencies among $(\omega_1, \dots, \omega_N)$ that are positive, so that $C_{N,+}$ captures the lack of symmetry of the initial chosen disorder: if $\alpha_N = 0$, there are as many positive frequencies as negative frequencies[2], whereas the case $\alpha_N > 0$ (resp. $\alpha_N < 0$) represents the case of an asymmetry in favor of positive (resp. negative) frequencies.

Numerical simulations (Figure 3) of the process given by (1.7) show a non self-averaging phenomenon for the fluctuation process compatible with the one observed for the rotation of the finite system (Figure 2): as seen in Figure 3, the trajectories of $\eta(\sin)$ have an approximately linear behavior, which slope depend on the realization of the initial mean-value C. Hence, a way to understand the dependence of the microscopic system (1.4) in the initial asymmetry of the disorder is to analyze the dependence of the fluctuation process η (1.7) in the mean-value C of its initial condition. More explicitly, the key point of this paper is to understand how different initial conditions in evolution (1.7) may lead to distinct approximately linear trajectories of the fluctuation process.

We tackle this issue via a spectral analysis of the evolution operator L_{q_t} defined in (1.8) via arguments from the general theory of perturbations of self-adjoint operators ([23]) and analytic semigroup of operators ([27, 33]) and usual techniques about SPDEs in Hilbert spaces ([9]), using the precise knowledge we have about the same linear operator L_{q_0} in the case without disorder (see [5]).

---

[2] in such case, the number N of rotators is necessarily even.



FIGURE 3. We plot here the evolution of the process $\eta_t(\sin)$, for different realizations of C; the trajectories are sample-dependent and compatible with the ones observed in Fig 2.

1.4. **Organization of the paper.** The paper is organized as follows: in Section 2, we precise the main set-up for the study of the SPDE (1.7) and state the main results. In particular, Theorem 2.5 and Theorem 2.7 deal with the spectral properties of the evolution operator L_{q_t} at least when the disorder is small. Secondly, we state the main result of this paper: Theorem 2.9 establishes the linear asymptotics of the fluctuation process solution of (1.7).

Section 3 is devoted to prove Theorem 2.5. In Section 4 we prove Theorem 2.7, whereas the main result of the paper, Theorem 2.9 is proved in Section 5.

2. MAIN DEFINITIONS AND RESULTS

In what follows, the following standard notations will be used: for an operator F, we will denote by $\rho(F)$ the set of all complex numbers λ for which $\lambda - F$ is invertible, and by $R(\lambda, F) := (\lambda - F)^{-1}$, $\lambda \in \rho(F)$ the resolvent of F. The spectrum of F will be denoted as $\sigma(F)$.

2.1. **The evolution operator L_q.** The main issue of this paper is to prove asymptotic results for the solution of the SPDE (1.7) as $t \to +\infty$; this fact is deeply linked to the spectral properties of the operator L_q defined in (1.8). We will restrict ourselves to the stationary case, that is when $q|_{t=0} = q_t$ is equal to the synchronized (non-trivial) stationary solution q (1.13) of evolution (1.1). In this case, the object of interest is the stationary version of (1.8), that is

$$Lh(\theta,\omega) := \frac{1}{2}\partial_\theta^2 h(\theta,\omega) - \partial_\theta\left(h(\theta,\omega)\left(\langle J * q\rangle_\mu(\theta) + \omega\right) + q(\theta,\omega)\langle J * h\rangle_\mu(\theta)\right) . \tag{2.1}$$

The domain $\mathcal{D}$ of the operator L is given by:

$$\mathcal{D} := \left\{h(\theta,\omega);\ \forall\omega, \theta \mapsto h(\theta,\omega) \in \mathcal{C}^2(\mathbf{S}),\ \int_{\mathbf{S}\times\mathbf{R}} h(\theta,\omega)\,\mathrm{d}\theta\mu(\,\mathrm{d}\omega) = 0\right\}. \tag{2.2}$$

Remark 2.1. *We point out that the choice of the domain $\mathcal{D}$ of L is of prime importance for our study of evolution* (1.7)*. In another context, one encounters the same operator L in the study of the linear stability of the stationary solution q since the linearized evolution of* (1.1) *around q is precisely given by $\partial_t h_t = Lh_t$. The natural domain for this latter evolution (see* [17]*) is then*

$$\left\{h(\theta,\omega);\ \forall\omega, \theta \mapsto h(\theta,\omega) \in \mathcal{C}^2(\mathbf{S}),\ \forall\omega,\ \int_{\mathbf{S}} h(\theta,\omega)\,\mathrm{d}\theta = 0\right\}. \tag{2.3}$$

Indeed for all ω, $q(\cdot,\omega)$ is a probability density on $\mathbf{S}$ so that perturbing by elements of domain (2.3) *enables to remain within the set of functions with integral* 1 *on* $\mathbf{S}$.

But here, the key point is to understand that evolution (1.7) *does not live in domain* (2.3) *since the fluctuations of the disorder are precisely captured by the fact that η has a non-trivial mean-value $C(\omega)$ for fixed ω. We will henceforth work with the domain $\mathcal{D}$ defined in* (2.2) *instead of* (2.3)*. This choice of domain (apart from being technically more demanding) is all the more relevant since we will see that the non self-averaging phenomenon holds in* (2.2) *and not in* (2.3) *(see Remark 2.6 below).*

For the rest of this paper, we fix $K > 1$ and we restrict ourselves to the case where

$$\mu = \frac{1}{2}\left(\delta_{-\omega_0} + \delta_{\omega_0}\right), \tag{2.4}$$

where $\omega_0 > 0$ is a fixed parameter. This assumption (2.4) on μ appears to be quite restrictive, but generalizing parts of the results we present here to more general distributions μ does not seem to be straightforward. We refer to § 2.7 for a discussion on this topic.

The first main goal of this paper is to state a spectral decomposition of the operator L defined in (2.1), based on perturbation arguments from the non-disordered case $\mu = \delta_0$ (see § 1.3.2 and § 2.3).

2.2. Distribution spaces.

The spectral analysis of the operator L will be mostly carried out in spaces of distribution that have H^{-1} regularity w.r.t. θ. But the precise study of L requires to introduce weighted version of H^{-1} that we define here. We first focus on weighted-Sobolev spaces of functions $\theta \mapsto h(\theta)$ on $\mathbf{S}$ (§ 2.2.1) and then introduce the corresponding spaces for functions with disorder $(\theta,\omega) \mapsto h(\theta,\omega)$ on $\mathbf{S} \times \mathrm{Supp}(\mu)$ (§ 2.2.2):

2.2.1. *Weighted Sobolev spaces:* For any bounded positive weight function $k(\cdot)$ on $\mathbf{S}$ such that $\int_{\mathbf{S}} k(\theta)\,\mathrm{d}\theta = 1$, we may consider the space $\mathbf{L}^2_k$ closure of $\mathcal{C}(\mathbf{S})$ w.r.t. the norm:

$$\| h \|_{2,k} := \left(\int_{\mathbf{S}} \frac{h^2(\theta)}{k(\theta)}\,\mathrm{d}\theta\right)^{\frac{1}{2}}. \tag{2.5}$$

The decomposition of h into the sum of $\mathrm{Span}(k)$ and its orthogonal supplementary in $\mathbf{L}^2_k$ may be written as:

$$h = \left(\int_{\mathbf{S}} h\right)\cdot k + h_0, \tag{2.6}$$

where $\int_{\mathbf{S}} h_0 = 0$. Since h_0 is with zero mean value, each of its primitives are 2π-periodic. In particular, we can consider $\mathbf{H}_k^{-1}$ the closure of $\mathcal{C}(\mathbf{S})$ with respect to the following weighted Sobolev norm:

$$\|h\|_{-1,k} := \left(\left(\int_{\mathbf{S}} h\right)^2 + \int_{\mathbf{S}} \frac{\mathcal{H}_0^2}{k}\right)^{\frac{1}{2}}, \tag{2.7}$$

where $\mathcal{H}_0$ is the primitive of h_0 on $\mathbf{S}$ such that $\int_{\mathbf{S}} \frac{\mathcal{H}_0}{k} = 0$. Note that one can understand the spaces $\mathbf{H}_k^{-1}$ as part of a *Gelfand-triple* construction (see Appendix A for a precise definition). In particular, we will make a constant use of the space $\mathbf{H}_{q_0}^{-1}$ (that is for $k(\cdot) = q_0(\cdot)$ where q_0 is the stationary solution (1.18) of the non-disordered system) which is the natural space (see Prop. 2.4) for the study on the Kuramoto operator L_{q_0} (2.11) in the non-disordered case.

Remark 2.2. *In the case of a constant weight* $k(\cdot) \equiv \frac{1}{2\pi}$, *we will write* $(\mathbf{L}^2, \|\cdot\|_2)$ *and* $(\mathbf{H}^{-1}, \|\cdot\|_{-1})$ *instead of* $(\mathbf{L}^2_{\frac{1}{2\pi}}, \|\cdot\|_{2,\frac{1}{2\pi}})$ *and* $(\mathbf{H}^{-1}_{\frac{1}{2\pi}}, \|\cdot\|_{-1,\frac{1}{2\pi}})$.

2.2.2. *Weighted Sobolev spaces (with disorder):* The natural space in where to study the operator L is the space of functions h in $\mathcal{D}$ such that each component $h(\cdot,\omega)$ live in a certain $\mathbf{H}^{-1}_{k(\cdot,\omega)}$ for a weight $k(\cdot,\omega)$ (which may depends on $\omega \in \mathrm{Supp}(\mu)$). More precisely, for any family of positive weight functions $(k(\cdot,\omega))_{\omega\in\mathrm{Supp}(\mu)}$, we denote as $\mathbf{H}^{-1}_{\mu,k}$ the closure of $\mathcal{D}$ w.r.t. the norm:

$$\|h\|_{\mu,-1,k} := \left(\int_{\mathbf{R}} \|h(\cdot,\omega)\|^2_{-1,k(\cdot,\omega)}\mu(\,\mathrm{d}\omega)\right)^{\frac{1}{2}} = \left(\int_{\mathbf{R}}\left(\int_{\mathbf{S}} h\,\mathrm{d}\theta\right)^2 \mathrm{d}\mu + \int_{\mathbf{R}}\int_{\mathbf{S}} \frac{\mathcal{H}_0^2}{k}\,\mathrm{d}\theta\,\mathrm{d}\mu\right)^{\frac{1}{2}}. \tag{2.8}$$

We will also consider the analogous averaged weighted $\mathbf{L}^2$-spaces, that is, for any family of positive weights $(k(\cdot,\omega))_{\omega\in\mathrm{Supp}(\mu)}$, the space $\mathbf{L}^2_{\mu,k}$ given by the norm:

$$\|h\|_{\mu,2,k} := \left(\int_{\mathbf{R}}\int_{\mathbf{S}} \frac{h(\theta,\omega)^2}{k(\theta,\omega)}\,\mathrm{d}\theta\,\mathrm{d}\mu(\omega)\right)^{\frac{1}{2}}. \tag{2.9}$$

Remark 2.3. *In the particular case of* $k(\cdot,\omega) \equiv \frac{1}{2\pi}$ *for all* $\omega \in \mathrm{Supp}(\mu)$, *we will write*

$$\mathbf{H}^{-1}_{\mu} := \mathbf{H}^{-1}_{\mu,\frac{1}{2\pi}}$$

and the corresponding norm will be denoted as $\|\cdot\|_{\mathbf{H}_\mu}$. *We will also write* $(\mathbf{L}^2_\mu, \|\cdot\|_{\mu,2})$ *instead of* $(\mathbf{L}^2_{\mu,\frac{1}{2\pi}}, \|\cdot\|_{\mu,2,\frac{1}{2\pi}})$.

The main theorem concerning the operator L (Theorem 2.7) will be stated in $\mathbf{H}^{-1}_\mu$ for the ease of exposition but its proof will require the introduction of weighted-Sobolev spaces $\mathbf{H}^{-1}_{\mu,k}$ for nontrivial weights k.

2.3. **The non-disordered case.** In the context of the Kuramoto model without disorder, the linearized operator L_{q_0} around stationary solution q_0 (see § 1.3.2), with domain

$$\mathcal{D}_0 := \left\{u \in \mathcal{C}^2(\mathbf{S});\ \int_{\mathbf{S}} u = 0\right\} \tag{2.10}$$

is:

$$L_{q_0}u := \frac{1}{2}\partial_\theta^2 u - \partial_\theta\left[q_0(J*u) + u(J*q_0)\right]. \tag{2.11}$$

In [5], it is mainly proved that L_{q_0} is essentially self-adjoint in $\mathbf{H}^{-1}_{q_0}$:

Proposition 2.4 (Bertini-Giacomin-Pakdaman). *$(L_{q_0}, \mathcal{D}_0)$ is essentially self-adjoint in $\mathbf{H}^{-1}_{q_0}$. The spectrum of (the self-adjoint extension of) L_{q_0} is pure point lying in $(-\infty, 0)$; 0 is in the spectrum, with one-dimensional eigenspace (spanned by $\partial_\theta q_0$). Moreover, the distance $\lambda_K(L_{q_0})$ between the eigenvalue 0 and the rest of the spectrum is strictly positive.*

Proposition 2.4 is crucial for our study of operator L (2.1). We will refer to it when required.

2.4. **Non self-averaging phenomenon for the operator L and existence of a Jordan block.** The linear trajectories that depend on the initial condition as observed in Figure 3 are reminiscent of an analogous deterministic finite-dimensional example: let us consider the 2-dimensional evolution $\left(\begin{smallmatrix} x'(t) \\ y'(t) \end{smallmatrix}\right) = L \left(\begin{smallmatrix} x(t) \\ y(t) \end{smallmatrix}\right)$, for $L = \left(\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}\right)$. Then it is trivial to see that the solutions of this system are linear in time: $\frac{x(t)}{t} \to y_0$ as $t \to \infty$. This can be easily generalized for matrices L $n \times n$ with spectrum in $\{\lambda \in \mathbf{C};\, \Re(\lambda) \leqslant 0\}$ such that the Jordan decomposition of L admits a $\left(\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}\right)$-block; this fact is precisely equivalent to the existence of x and y such that $Lx = 0$ and $Ly = x$. The purpose of the first main theorem of this paper is to prove an analogous existence of a Jordan block for the operator L:

Theorem 2.5. *For any fixed $\omega_0 > 0$, if q is the stationary solution in* (1.13)*, then*

$$L \partial_\theta q = 0. \tag{2.12}$$

Moreover, there exists $p \in \mathcal{D}$ such that

$$\forall \theta \in \mathbf{S}, \forall \omega \in \operatorname{Supp}(\mu), \qquad Lp(\theta, \omega) = \partial_\theta q(\theta, \omega). \tag{2.13}$$

In particular, the characteristic space of L in 0 *is* at least *of dimension* 2*.*

Remark 2.6. *Equality* (2.12) *is a direct consequence of the rotation invariance in* (1.4) *(Remark 1.1). Note also that $p(\cdot, \omega)$ found in* (2.13) *is with non-trivial mean value for all $\omega \in \operatorname{Supp}(\mu)$. Indeed, we believe that $\int_{\mathbf{S}} p(\cdot, \omega)$ is in fact equal to $-\frac{1}{\omega}$; this fact is derived from non-rigorous computations and verified by numerical simulations. In other terms, if we had chosen domain* (2.3) *instead of domain* (2.2)*, we would not have found such p; the Jordan block $\left(\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}\right)$ in the matrix representation* (2.17) *of the operator L does not exist on the domain* (2.3)*.*

Theorem 2.5 is proved in Section 3.

2.5. **Spectral properties of L and position of the spectrum.** The second goal of this paper is to prove that L generates an analytic semi-group of operators with spectrum confined in the part of the complex plane with negative real part:

Theorem 2.7. *In the Hilbert space $\mathbf{H}^{-1}_{\mu}$ defined in Remark 2.3, the operator $(L, \mathcal{D})$ is densely defined, closable, its closed extension having compact resolvent. In particular, its spectrum consists of isolated eigenvalues with finite multiplicities.*

Moreover, for all $K > 1$, for all $\alpha \in (0, \frac{\pi}{2})$, for all $\rho \in (0, 1)$, there exists $\omega_\star = \omega_\star(K, \alpha, \rho) > 0$ such that, for all $0 < \omega_0 < \omega_\star$, the following is true:

- *The spectrum of L lies in a cone C_α with vertex* 0 *and angle α*

$$C_\alpha := \left\{\lambda \in \mathbb{C};\, \frac{\pi}{2} + \alpha \leqslant \arg(\lambda) \leqslant \frac{3\pi}{2} - \alpha\right\} \subseteq \{z \in \mathbf{C};\, \Re(z) \leqslant 0\}\ ; \tag{2.14}$$

- *There exists $\alpha' \in (0, \frac{\pi}{2})$ such that L is the infinitesimal generator of an analytic semi-group defined on a sector $\Delta_{\alpha'} := \{\lambda \in \mathbb{C},\, |\arg(\lambda)| < \alpha'\}$;*

- *the dimension of the characteristic space in* 0 *is* exactly 2, *spanned by* $\partial_\theta q$ *and* p, *where* p *is defined in Theorem 2.5,*
- *the eigenvalue* 0 *is separated from the rest of the spectrum at a distance* $\lambda_K(L) = \lambda(L, K, \rho)$ *at least equal to* $\rho \cdot \min\left(\lambda_K(L_{q_0}), \frac{1}{2}e^{-4Kr_0}\right)$, *where* L_{q_0} *and* r_0 *are defined in § 1.3.2.*

Note that Theorem 2.7 relies on perturbation arguments of the non-disordered case mentioned in § 2.3; in that sense, the spectral gap $\lambda_K(L)$ found in Theorem 2.7 depends on the spectral gap $\lambda_K(L_{q_0})$ for the non-disordered case.

As a consequence of Theorem 2.7, there exists a decomposition of the space $\mathbf{H}_\mu^{-1}$ into the direct sum

$$\mathbf{H}_\mu^{-1} = G_0 \oplus G_{<0}\,, \tag{2.15}$$

where G_0 is of dimension 2 (spanned by $\partial_\theta q$ and p) such that the restriction of the operator L to G_0 has spectrum $\{0\}$ and the restriction of L to $G_{<0}$ has spectrum $\sigma(L) \smallsetminus \{0\} \subseteq \{\lambda \in \mathbf{C};\, \Re(\lambda) < 0\}$. We will denote as P_0 the corresponding projection on G_0 along to $G_{<0}$, and $P_{<0} = 1 - P_0$. In particular, there exist unique continuous linear forms $\ell_{\partial_\theta q}$ and ℓ_p such that for all $h \in \mathbf{H}_\mu^{-1}$

$$P_0 h := \ell_{\partial_\theta q}(h)\partial_\theta q + \ell_p(h)p. \tag{2.16}$$

To fix ideas, one may think of the following infinite matrix representation for the operator L:

$$L = \begin{array}{c} \left(\begin{array}{c|c} P_0\, L P_0 & P_0\, L P_{<0} \\ \hline P_{<0}\, L P_0 & P_{<0}\, L P_{<0} \end{array}\right) \\ \begin{array}{cc} \underbrace{\qquad\qquad}_{\substack{G_0 = \\ \mathrm{Span}(\partial_\theta q, p)}} & \underbrace{\qquad\qquad}_{G_{<0}} \end{array} \end{array} = \begin{array}{c} \left(\begin{array}{cc|ccc} 0 & 1 & \multicolumn{3}{c}{\ell_{\partial_\theta q}(LP_{<0})} \\ 0 & 0 & 0 & \cdots & 0 \\ \hline 0 & 0 & & & \\ \vdots & \vdots & & P_{<0}\, L P_{<0} & \\ 0 & 0 & & & \end{array}\right) \\ \begin{array}{ccc} \underbrace{\ }_{\partial_\theta q} & \underbrace{\ }_{p} & \underbrace{\qquad\qquad}_{G_{<0}} \end{array} \end{array} \begin{array}{l} \left.\begin{array}{l} \}\partial_\theta q \\ \}p \end{array}\right\} G_0 \\ \left.\begin{array}{l} \\ \\ \\ \end{array}\right\} G_{<0} \\ \\ \end{array} \tag{2.17}$$

Note that the second line in the matrix representation (2.17) of L is indeed equally zero since for all $h \in \mathbf{H}_\mu^{-1}$, Lh is of zero mean value on $\mathbf{S}$; in particular $\ell_p(Lh) = 0$ for all $h \in \mathbf{H}_\mu^{-1}$.

Remark 2.8. *Any element* $h = (h(\theta, \omega))_{\theta \in \mathbf{S}, \omega \in \mathrm{Supp}(\mu)}$ *can be identified in our binary case* (2.4) *with a couple* $(h_+(\theta), h_-(\theta))_{\theta \in \mathbf{S}}$. *Moreover, any* $h \in \mathbf{H}_\mu^{-1}$ *can be decomposed according to* (2.15)*:*

$$h = \ell_{\partial_\theta q}(h)\partial_\theta q + \ell_p(h)p + P_{<0}\, h.$$

Let us integrate the latter decomposition w.r.t. θ. *Since* $\int_{\mathbf{S}} Lu = 0$ *for all* $u \in \mathcal{D}$, *we have* $\int_{\mathbf{S}} P_{<0}\, h = 0$ *so that one can actually find an explicit formulation for the functional* ℓ_p*:*

$$\ell_p(h) = \frac{\int_{\mathbf{S}} h_+}{\int_{\mathbf{S}} p_+} = \frac{\int_{\mathbf{S}} h_-}{\int_{\mathbf{S}} p_-}. \tag{2.18}$$

The last equality in (2.18) *is due to the fact that* $\int_{\mathbf{S}} (h_+ + h_-) = \int_{\mathbf{S}} (p_+ + p_-) = 0$.

2.6. **Long time evolution of the fluctuation SPDE.** We now turn to the main result of the paper, which concerns the asymptotic behavior of the fluctuation process η defined in (1.7):

Theorem 2.9. *Under the hypothesis of Theorem 2.7, there exists a unique weak solution η to* (1.7) *in $\mathbf{H}_\mu^{-1}$. Moreover, η satisfies the following asymptotic linear behavior: for fixed initial condition $\eta_0^\omega = X + C(\omega)$*

$$\frac{\eta_t}{t} \xrightarrow[t\to\infty]{\text{in law}} \frac{\int_{\mathbf{S}} C_+(\omega)}{\int_{\mathbf{S}} p_+} \partial_\theta q := v(\omega)\partial_\theta q, \quad \text{as } t \to +\infty, \tag{2.19}$$

where $p_+(\theta) := p(\theta, \omega_0)$ is defined by (2.13)*. Moreover, in view of the structure of the initial condition η_0 (recall* (1.23)*), the speed $v(\cdot)$ is a Gaussian random variable in ω with variance*

$$\sigma_v^2 := \left(2\int_{\mathbf{S}} p_+(\theta)\,\mathrm{d}\theta\right)^{-2}. \tag{2.20}$$

Hence, Theorem 2.9 gives an explicit expression of the speed of rotation of the fluctuation process.

2.7. Perspectives.

2.7.1. *Possible generalizations.* In [4, § 10.2, p. 47], Balmforth and Sassi found on the basis of numerical experiments that if we get rid artificially of the asymmetry between positive frequencies and negative frequencies, the disorder-induced rotation in (1.4) disappears. More precisely, the simulations carried out in [4] are ruled by the following principle: consider a sample of $2N$ rotators ($N \geqslant 1$) and pick up at random the frequencies ω_i for exactly N rotators. For each of those N frequencies ω_i, attribute to one the remaining N rotators the exact opposite frequency $-\omega_i$. The conclusion of [4] is that for this choice of frequencies, there is no rotation, no matter how the first N frequencies are sampled. We stress the fact that we retrieve this phenomenon in Theorem 2.9 in the case where the first N rotators are i.i.d. samples of $\mu = \frac{1}{2}(\delta_{-\omega_0} + \delta_{\omega_0})$, since in that case the quantity α_{2N} defined in (1.26) is equally zero for all $N \geqslant 1$ and the consequent limit speed v is equally zero in this case.

One could hope to generalize the results of the paper in at least two directions. Firstly, we restrict ourselves to the binary case $\mu = \frac{1}{2}(\delta_{-\omega_0} + \delta_{\omega_0})$. Note that the proof of Theorem 2.5 concerning the existence of a Jordan block (although written in this particular case for the reader's convenience) is not specific to this restrictive case: one could easily rewrite the same proof in the case of more general distributions μ (even with unbounded support), satisfying appropriate integrability conditions in 0 and in ∞.

The main restriction on μ concerns Theorem 2.7: the hypothesis $\mu = \frac{1}{2}(\delta_{-\omega_0} + \delta_{\omega_0})$ is critical for its proof. In that sense, there are some deeper considerations which may prove possible generalizations of Theorem 2.7 to be difficult: indeed, the key argument of the proof is based on the fact that perturbing a finite dimensional kernel of an operator A by a sufficiently small perturbation B leads to a kernel for the operator $A + B$ with the same finite dimension. But for distributions more general than binary distributions $\frac{1}{2}(\delta_{-\omega_0} + \delta_{\omega_0})$, the kernel of L is likely to become of infinite dimension, so that those perturbation arguments cannot be applied.

Secondly, Theorem 2.7 is only proved for small disorder ω_0 whereas one would expect it to be true even for large disorder. It is indeed natural to believe that the non self-averaging phenomenon seen in Figure 2 not only holds for large disorder but would even be more noticeable in that case. However, since Theorem 2.7 relies on perturbation arguments, to prove similar results for large ω_0 would seem to require alternative methods.

2.7.2. *Other possible time scales.* More importantly, Theorem 2.9 gives an expression of the speed at which the limit of the fluctuation process η rotates. Since in this work, we are only considering time scales of order $O(1)$ (that is for arbitrary but fixed time horizon T), this may not be the right speed of rotation for the finite-size system: on larger time-scales (or order $O(\sqrt{N})$ say), there may be accumulation of fluctuations which may lead to a different expression of the speed of rotation. Hence, to understand this phenomenon properly, an interesting result would be the study of the rescaled empirical process $(\nu_{\sqrt{N}t})_{t,N}$, as N and t are large. This has not been carried out in this article and would be a natural perspective for this work.

3. On the existence of a Jordan block for L (Proof of Theorem 2.5)

The purpose of this section is to prove Theorem 2.5, that is the fact that the operator L defined in (2.1) has a Jordan block of size at least 2. The symmetry of the system (recall Remark 1.1) leads to consider the set of distributions which are odd w.r.t. the variable $(\theta,\omega) \in \mathbf{S} \times \mathrm{Supp}(\mu)$:

$$\mathcal{O} := \{h;\ \forall(\theta,\omega) \in \mathbf{S} \times \mathrm{Supp}(\mu),\ h(-\theta,-\omega) = -h(\theta,\omega)\}\,. \tag{3.1}$$

We also denote by $\mathcal{N}$ the set of functions that with zero mean-value for all $\omega \in \mathrm{Supp}(\mu)$ (recall the definition of $\mathcal{D}$ in (2.2)):

$$\mathcal{N} := \left\{h \in \mathcal{D};\ \forall\omega \in \mathrm{Supp}(\mu),\ \int_{\mathbf{S}} h(\theta,\omega)\,\mathrm{d}\theta = 0\right\}. \tag{3.2}$$

In the following straightforward lemma, whose proof is left to the reader, we sum-up the basic properties of the stationary solution q (recall (1.13)):

Lemma 3.1. *The following statements are true:*

1. $\partial_\theta q \in \mathcal{O}$,
2. *If* $h \in \mathcal{O}$ *then* $Lh \in \mathcal{O}$,
3. $\partial_\theta q \in \mathcal{N}$, *and for all* $h \in \mathcal{D}$, $Lh \in \mathcal{N}$,
4. *There exist some positive constants* c *and* C *such that for all* $\theta \in \mathbf{S}$, $\omega \in \mathrm{Supp}(\mu)$, $0 < c \leqslant q(\theta,\omega) \leqslant C$,
5. *For all* $\theta \in \mathbf{S}$, $\omega \in \mathrm{Supp}(\mu)$,
$$\frac{1}{2}\partial_\theta q(\theta,\omega) = q(\theta,\omega)\left(\langle J * q\rangle_\mu + \omega\right) + \kappa(\omega)\,, \tag{3.3}$$
where $\kappa(\omega) = \frac{1-e^{4\pi\omega}}{2Z(\omega)}$, *and* $Z(\omega) = Z(\omega, 2Kr)$ *is the constant of normalization defined in* (1.13).

Note in particular that the fact that $\partial_\theta q \in \mathcal{O}$ can be seen as a consequence of Remark 1.1. A direct calculation easily shows that $\partial_\theta q$ is in the kernel of L (it corresponds to the rotation invariance of the problem). The rest of this section is devoted to prove the existence of an element $p \in \mathcal{D}$ such that $Lp = \partial_\theta q$.

We recall here the definition of the weighted Sobolev spaces introduced in § 2.2.2: we make use here of the space $\left(\mathbf{H}^{-1}_{\mu,q}, \langle\cdot\,,\,\cdot\rangle_{\mu,-1,q}\right)$ defined in (2.8) in the case of $k = q$ and of the space $\left(\mathbf{L}^2_\mu, \langle\cdot\,,\,\cdot\rangle_{\mu,2}\right)$ defined in Remark 2.3. The main result is the following:

Proposition 3.2. *For every* $\omega_0 > 0$, *in the binary case* (2.4), *for every* $v \in \mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O}$ *(and in particular for* $v = \partial_\theta q$*), there exists some* $p \in \mathbf{L}^2_\mu \cap \mathcal{O}$ *such that*

$$\forall l \in \mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O},\ \langle Lp\,,\,l\rangle_{\mu,-1,q} = \langle v\,,\,l\rangle_{\mu,-1,q}\,. \tag{3.4}$$

Moreover, in the case $v = \partial_\theta q$*, any* p *that satisfies* (3.4) *is in fact a regular function* ($p(\cdot,\omega) \in \mathcal{C}^\infty(\mathbf{S})$ *for all* $\omega \in \mathrm{Supp}(\mu)$) *and is a classical solution to* (2.13)*.*

Remark 3.3. *The scope of Proposition 3.2 is more general than the restrictive case of a binary law* $\mu = \frac{1}{2}\left(\delta_{-\omega_0} + \delta_{\omega_0}\right)$*; indeed the following proof works for more general distributions* μ*, the only additional requirement being integrability conditions*[3] *in* 0 *and* $+\infty$*, see Remark 3.9.*

Proof of Proposition 3.2 relies on several lemmas:

Lemma 3.4. *For* $h \in \mathcal{O} \cap \mathcal{D}$*,* $l \in \mathcal{O} \cap \mathcal{D}$*, let us introduce the Dirichlet form*

$$\mathcal{E}_L(h,l) := \langle Lh\,,\, l\rangle_{\mu,-1,q}\,. \tag{3.5}$$

$\mathcal{E}_L(\cdot,\cdot)$ *is well defined on* $\mathcal{D}(\mathcal{E}_L) := \left(\mathbf{L}^2_\mu \cap \mathcal{O}\right) \times \left(\mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O}\right)$ *and one can decompose* $\mathcal{E}_L(\cdot,\cdot)$ *into:*

$$\forall (h,l) \in \mathcal{D}(\mathcal{E}_L),\ \mathcal{E}_L(h,l) = \Gamma(h,l) + K\ell(h)\cdot \ell(l), \tag{3.6}$$

where $\Gamma(\cdot,\cdot)$*, bilinear form on* $\mathcal{D}(\mathcal{E}_L)$ *and* $\ell(\cdot)$ *linear form on* $\mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O}$*, are defined as follows:*

$$\forall (h,l) \in \mathcal{D}(\mathcal{E}_L),\ \Gamma(h,l) := -\frac{1}{2}\int_{\mathbf{S}\times\mathbf{R}} \frac{hl}{q}\,\mathrm{d}\lambda\,\mathrm{d}\mu + \int_{\mathbf{S}\times\mathbf{R}} \kappa(\cdot)\frac{h\mathcal{L}}{q^2}\,\mathrm{d}\lambda\,\mathrm{d}\mu, \tag{3.7}$$

$$\forall l \in \mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O},\ \ell(l) := \int_{\mathbf{S}\times\mathbf{R}} l\sin(\cdot)\,\mathrm{d}\lambda\,\mathrm{d}\mu, \tag{3.8}$$

where κ *and* $\mathcal{L}$ *in* (3.7) *are respectively defined in* (3.3) *and as the primitive of* $l \in \mathbf{H}^{-1}_{\mu,q}$ *such that* $\int_{\mathbf{S}} \frac{\mathcal{L}(\cdot,\omega)}{q(\cdot,\omega)} = 0$ *for all* $\omega \in \mathrm{Supp}(\mu)$ *(recall § 2.2.2).*

Lemma 3.5. *For all continuous linear form* f *on* $\mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O}$*, there exists some* $p_1 \in \mathbf{L}^2_\mu \cap \mathcal{O}$ *such that for* $l \in \mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O}$

$$\Gamma(p_1,l) = f(l). \tag{3.9}$$

Lemma 3.6. *The linear form* $\ell(\cdot)$ *defined in* (3.8) *can be expressed as a scalar product on* $\mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O}$*: there exists* $p_2 \in \mathcal{D} \cap \mathcal{O}$*, for all* $l \in \mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O}$

$$\ell(l) = \langle Lp_2\,,\, l\rangle_{\mu,-1,q}\,. \tag{3.10}$$

Let us admit for a moment Lemmas 3.4, 3.5 and 3.6 and let us prove Proposition 3.2:

Proof of Proposition 3.2. Let v be a fixed element of $\mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O}$. Applying Lemma 3.5 to the continuous linear form $f(l) = \langle v\,,\, l\rangle_{\mu,-1,q}$, there exists some $p_1 \in \mathbf{L}^2_\mu \cap \mathcal{O}$ such that $\Gamma(p_1,l) = \langle v\,,\, l\rangle_{\mu,-1,q}$, which gives using Lemma 3.4 and Lemma 3.6:

$$\begin{aligned}\langle v\,,\, l\rangle_{\mu,-1,q} &= \Gamma(p_1,l),\\ &= \langle Lp_1\,,\, l\rangle_{\mu,-1,q} - K\ell(p_1)\ell(l),\\ &= \langle Lp_1\,,\, l\rangle_{\mu,-1,q} - K\ell(p_1)\langle Lp_2\,,\, l\rangle_{\mu,-1,q}\,.\end{aligned}$$

We can conclude that the variational formula (3.4) is verified for the following choice of p:

$$p := p_1 - K\ell(p_1)p_2 \in \mathbf{L}^2_\mu \cap \mathcal{O}. \tag{3.11}$$

Let us prove now that such p is in fact a regular function in θ: since $p_2 \in \mathcal{D}$ is regular in θ, it suffices to prove that for all $\omega \in \mathrm{Supp}(\mu)$, $\theta \mapsto p_1(\theta,\omega)$ is $\mathcal{C}^2$ (in fact $\mathcal{C}^\infty$) in θ. We start from the definition of p_1:

$$\forall l \in \mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O}, \quad \Gamma(p_1,l) = \langle \partial_\theta q\,,\, l\rangle_{\mu,-1,q}\,.$$

---

[3] those conditions are obviously satisfied in the binary case (2.4).

Since this true for all $l \in \mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O}$, thanks to the expression of Γ in (3.7), we obtain that for any fixed $\omega \in \mathrm{Supp}(\mu)$, for Lebesgue-almost every $\theta \in \mathbf{S}$:

$$\frac{1}{4}\frac{p_1(\theta,\omega)}{q(\theta,\omega)} = -\kappa(\omega)\left(\int_0^\theta \frac{p_1(u,\omega)}{q(u,\omega)^2}\,\mathrm{d}u\right) + \int_0^\theta \frac{Q(u,\omega)}{q(u,\omega)}\,\mathrm{d}u, \tag{3.12}$$

where $Q(\cdot,\omega)$ is the primitive of $\partial_\theta q(\cdot,\omega)$ such that $\int_{\mathbf{S}} \frac{Q(\cdot,\omega)}{q(\cdot,\omega)} = 0$. But, using that q is bounded and $\mathcal{C}^\infty$ in θ and that $p_1(\cdot,\omega) \in \mathbf{L}^2$, the primitive $\int_0^\theta \frac{p_1(u,\omega)}{q(u,\omega)^2}\,\mathrm{d}u$ as a $\mathcal{C}^1$ version. So, thanks to (3.12), p_1 has a $\mathcal{C}^1$ version. So, the right-hand side of (3.12) is a least $\mathcal{C}^2$, and so does p_1. The same repeated argument shows that p_1 is $\mathcal{C}^\infty$ in θ. That concludes the proof of Proposition 3.2. □

It now remains to prove the three lemmas:

Proof of Lemma 3.4. Let us prove equality (3.6): since $\mathcal{L}$ is a primitive of l, one has

$$\frac{1}{2}\int_{\mathbf{S}} \frac{(\partial_\theta h)\mathcal{L}}{q} = -\frac{1}{2}\int_{\mathbf{S}} \frac{hl}{q} + \frac{1}{2}\int_{\mathbf{S}} \frac{h\mathcal{L}}{q^2}\partial_\theta q\,.$$

Using (3.3), for $\omega \in \mathrm{Supp}(\mu)$

$$\frac{1}{2}\int_{\mathbf{S}} \frac{(\partial_\theta h)\mathcal{L}}{q} = -\frac{1}{2}\int_{\mathbf{S}} \frac{hl}{q} + \int_{\mathbf{S}} \frac{h\mathcal{L}}{q}\left(\langle J * q\rangle_\mu(\cdot) + \omega\right) + \kappa(\omega)\int_{\mathbf{S}} \frac{h\mathcal{L}}{q^2}.$$

Thanks to the expression of Lh in (2.1), we obtain

$$\begin{aligned}\mathcal{E}_L(h,l) &= -\frac{1}{2}\int_{\mathbf{S}\times\mathbf{R}} \frac{hl}{q} + \int_{\mathbf{S}\times\mathbf{R}} \kappa(\cdot)\frac{h}{q^2}\mathcal{L} - \int_{\mathbf{S}\times\mathbf{R}} \langle J * h\rangle_\mu \mathcal{L}, \\ &= \Gamma(h,l) - \int_{\mathbf{S}\times\mathbf{R}} \langle J * h\rangle_\mu \mathcal{L}.\end{aligned} \tag{3.13}$$

Lastly, integrating by parts the last term in (3.13) and expanding the cosine function (recall $J(\cdot) = -K\sin(\cdot)$), we obtain:

$$\begin{aligned}-\int_{\mathbf{S}\times\mathbf{R}} \langle J * h\rangle_\mu \mathcal{L} = K&\left(\int_{\mathbf{S}\times\mathbf{R}} \cos(\cdot) l\,\mathrm{d}\lambda\,\mathrm{d}\mu\right)\left(\int_{\mathbf{S}\times\mathbf{R}} \cos(\cdot) h\,\mathrm{d}\lambda\,\mathrm{d}\mu\right) \\ &+ K\left(\int_{\mathbf{S}\times\mathbf{R}} \sin(\cdot) l\,\mathrm{d}\lambda\,\mathrm{d}\mu\right)\left(\int_{\mathbf{S}\times\mathbf{R}} \sin(\cdot) h\,\mathrm{d}\lambda\,\mathrm{d}\mu\right).\end{aligned}$$

But, since $l \in \mathcal{O}$, the first term in the latter expression is zero. The result (3.6) follows. □

Proof of Lemma 3.5. In this proof, we use the following extension to Lax-Milgram Theorem (cf. [38, chap. III]):

Proposition 3.7 (Lions-Lax-Milgram)**.** *Let $\{\mathcal{H}, |.|\}$ be a Hilbert space and $\{\mathcal{G}, \|\,.\,\|\}$ a normed linear space. Suppose $\Gamma : \mathcal{H}\times\mathcal{G} \to \mathbf{R}$ is bilinear and that $\Gamma(\cdot,\varphi)$ is continuous for each $\varphi \in \mathcal{G}$. If there exists some constant $C > 0$ such that*

$$\inf_{\|\varphi\|=1}\ \sup_{|h|\leqslant 1} |\Gamma(h,\varphi)| \geqslant C > 0, \quad \textit{(weak coercivity)}, \tag{3.14}$$

Then for each $f \in \mathcal{G}'$ there exists some $p \in \mathcal{H}$ such that $\Gamma(p,\varphi) = f(\varphi)$ for all $\varphi \in \mathcal{G}$.

The principle of the proof of Lemma 3.5 is to show that the bilinear function Γ defined in (3.7) satisfies Proposition 3.7 for

$$\mathcal{H} := \mathbf{L}^2_\mu \cap \mathcal{O}, \quad \text{endowed with} \quad \| \cdot \|_{\mu,2}\,, \tag{3.15}$$

$$\mathcal{G} := \mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O} \cap \mathbf{L}^\infty(\mathbf{S}\times\mathbf{R}), \quad \text{endowed with} \quad \| \cdot \|_{\mu,-1,q}\,. \tag{3.16}$$

Namely, we have the following:

(1) For each $l \in \mathcal{G}$, $\Gamma(\cdot, l)$is continuous on $\mathbf{L}^2_\mu \cap \mathcal{O}$: indeed, for the first term of $\Gamma(h,l)$, we have

$$\left| \int_{\mathbf{S}\times\mathbf{R}} \frac{hl}{q}\, \mathrm{d}\lambda\, \mathrm{d}\mu \right| \leqslant C \,\| \, l \,\|_\infty \int_{\mathbf{S}\times\mathbf{R}} |h|\, \mathrm{d}\lambda\, \mathrm{d}\mu \leqslant C \,\| \, h \,\|_{\mu,2}\,.$$

And for the second term, using the boundedness of q:

$$\int_{\mathbf{S}\times\mathbf{R}} \left| \kappa(\cdot) \frac{h}{q^2} \mathcal{L} \right| \, \mathrm{d}\lambda\, \mathrm{d}\mu \leqslant C \int_{\mathbf{S}\times\mathbf{R}} |h\mathcal{L}| \, \mathrm{d}\lambda\, \mathrm{d}\mu \leqslant C \,\| \, h \,\|_{\mu,2} \,\| \, l \,\|_{\mu,-1,q}\,.$$

(2) Γ is weakly coercive: let us fix $l \in \mathcal{G}$ such that $\| \, l \,\|_{\mu,-1,q} = 1$.

Let us choose $h = g\mathcal{L} \in \mathbf{L}^2_\mu \cap \mathcal{O}$, where for all $\omega \in \mathrm{Supp}(\mu)$, $g(\cdot,\omega)$ is a 2π-periodic function to be defined later. Then, by integration by parts in the equality (3.7)

$$\Gamma(h,l) = -\frac{1}{2} \int_{\mathbf{S}\times\mathbf{R}} \frac{g}{q} l \mathcal{L} + \int_{\mathbf{S}\times\mathbf{R}} \kappa(\cdot) \frac{g}{q^2} \mathcal{L}^2 = \int_{\mathbf{S}\times\mathbf{R}} \left\{ \frac{1}{4} \partial_\theta \left(\frac{g}{q} \right) + \kappa(\cdot) \frac{g}{q^2} \right\} \mathcal{L}^2\,. \tag{3.17}$$

Consider now for fixed $\omega \in \mathrm{Supp}(\mu)$ the following first order ODE, with periodic boundary condition:

$$\frac{1}{4} \partial_\theta f(\cdot,\omega) + \kappa(\omega) \frac{f(\cdot,\omega)}{q(\cdot,\omega)} = \frac{1}{q(\cdot,\omega)}, \quad \text{with } f(0,\omega) = f(2\pi,\omega). \tag{3.18}$$

Then for any $\omega \in \mathrm{Supp}(\mu) \setminus \{0\}$, an explicit calculation (left to the reader) shows that there exists a unique solution to (3.18), $\theta \mapsto f(\theta,\omega)$.

Remark 3.8. *In the case* $\omega = 0$, (3.18) *reduces to* $\frac{1}{4}\partial_\theta f = \frac{1}{q_0}$ *which is incompatible with the condition* $f(0) = f(2\pi)$, *since* $\int_{\mathbf{S}} \frac{1}{q_0}\, \mathrm{d}\theta > 0$*: there is no such* 2π*-periodic solution in the case* $\omega = 0$.

Moreover, it is straightforward to see that $\left\| \int_{\mathbf{R}} |f(\cdot,\omega)|\, \mathrm{d}\mu \right\|_{\infty,\mathbf{S}} \leqslant C$, for some constant $C > 0$.

Remark 3.9. *It is easy to see that* $f(\cdot,\omega)$ *is not bounded as* $\omega \to 0$ *and* $\omega \to +\infty$*; thus, for general distributions* μ*, the same control on* f *requires additional integrability assumptions in* 0 *and* $+\infty$ *(namely* $\int_{\mathbf{R}} \max\left(\frac{1}{|\omega|}, e^{c\omega} \right) \mu(\mathrm{d}\omega) < \infty$ *for some constant* $c > 0$*).*

If we choose h such that $h = g \cdot \mathcal{L}$ with $g(\cdot,\omega) = q(\cdot,\omega) f(\cdot,\omega)$, we have the following:

- By construction of f, using (3.18) in (3.17), $\Gamma(h,l) = \| \, l \,\|^2_{\mu,-1,q} = 1$,
- $\| \, h \,\|^2_{\mu,2} \leqslant C \int_{\mathbf{S}} f^2 \frac{\mathcal{L}^2}{q}\, \mathrm{d}\lambda \leqslant C$. So, $\sup_{\| \, h \,\|_{\mu,2,1} \leqslant 1} |\Gamma(h,l)| \geqslant \frac{1}{C}$, where C is independent of $l \in \mathcal{G}$ such that $\| \, l \,\|_{\mu,-1,q} = 1$.

Applying Proposition 3.7, we obtain the existence of some $p_1 \in \mathbf{L}^2_\mu \cap \mathcal{O}$ such that $\Gamma(p_1, l) = f(l)$, for all $l \in \mathcal{G}$. But by density, this is also true for $l \in \mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O}$. □

Proof of Lemma 3.6. Since there exists some constants $C, c > 0$ such that for all $\omega \in \mathrm{Supp}(\mu)$, $\theta \in \mathbf{S}$, $0 < c \leqslant q(\theta, \omega) \leqslant C$, ℓ is continuous on $\mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O}$ (as well as on $\mathbf{L}^2_\mu \cap \mathcal{O}$). More precisely, by Riesz theorem, there exists a unique $e \in \mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O}$ such that for all $l \in \mathbf{H}^{-1}_{\mu,q} \cap \mathcal{O}$, $\ell(l) = \langle e\,,\, l \rangle_{\mu,-1,q}$. One can be more explicit: a simple calculation shows that this $(\theta, \omega) \mapsto e(\theta, \omega)$ corresponds to the primitive $\mathcal{E}(\theta, \omega) = -q(\theta, \omega)\cos(\theta)$, that is:

$$\forall \theta \in \mathbf{S},\ \omega \in \mathrm{Supp}(\mu), \quad e(\theta, \omega) = -\partial_\theta q(\theta, \omega)\cos(\theta) + q(\theta, \omega)\sin(\theta). \tag{3.19}$$

Let us introduce the following function $p_2 \in \mathbf{L}^2_\mu \cap \mathcal{O}$:

$$p_2(\theta, \omega) = \frac{e^{-B(\theta,\omega)}}{1 - e^{4\pi\omega}} \int_{\mathbf{S}} e^{B(u,\omega) + 4\pi\omega}\,\mathrm{d}u + \int_0^\theta e^{B(u,\omega) - B(\theta,\omega)}\,\mathrm{d}u, \tag{3.20}$$

for

$$B(\theta, \omega) = -2\left(Kr\left(\cos(\theta) - 1\right) + \omega\theta\right).$$

Then one readily verifies that Lp_2 is proportional to e. ☐

4. GLOBAL SPECTRAL PROPERTIES OF OPERATOR L (PROOF OF THEOREM 2.7)

The purpose of this section is to prove Theorem 2.7. The main idea of the proof is to decompose the operator L defined by (2.1) on the domain $\mathcal{D}$ given by (2.2) into the sum of a self-adjoint operator A (in a weighted Sobolev space for appropriate weights) and a perturbation B which will be considered to be small w.r.t. A. Namely, it is straightforward to verify from (2.1) that L can be written as $L = A + B$ where, for all $h \in \mathcal{D}$, for all $\omega \in \mathrm{Supp}(\mu)$,

$$Ah(\theta, \omega) := \frac{1}{2}\partial_\theta^2 h(\theta, \omega) - \partial_\theta\Big(h(\theta, \omega)(J * q_0)(\theta) + q_0(\theta)\langle J * h \rangle_\mu\Big), \tag{4.1}$$

and,

$$Bh(\theta, \omega) := -\partial_\theta\Big(h(\theta, \omega)\{\langle J * (q - q_0)\rangle_\mu(\theta) + \omega\} + (q(\theta, \omega) - q_0(\theta))\langle J * h \rangle_\mu\Big). \tag{4.2}$$

We divide the proof of Theorem 2.7 into three parts: in § 4.1, we prove that A is essentially self-adjoint (and thus generates an analytic semigroup) in some weighted-Sobolev space (recall § 2.2) for an appropriate choice of weights. Note that this section strongly relies on the fact that μ is a binary distribution (recall (2.4)).

The purpose of § 4.2 is to establish precise control of the size of the perturbation B w.r.t. A. The third step of the proof, carried out in § 4.3, consists in deriving similar spectral properties for the operator $L = A + B$, especially the fact that the spectrum of L lies in the part of the complex plane with negative real part.

4.1. Spectral properties of the operator A.

In this paragraph, we prove mainly that A defined in (4.1) is essentially self-adjoint for a Sobolev norm that is equivalent to the norm $\|\cdot\|_{\mathbf{H}_\mu}$ defined in § 2.2.2.

Since we are working in the domain $\mathcal{D}$ (recall (2.2)), the test functions h are such that $\int_{\mathbf{R}} h(\cdot, \omega)\,\mathrm{d}\mu = \frac{1}{2}\left(h(\cdot, +\omega_0) + h(\cdot, -\omega_0)\right)$ has zero mean value on $\mathbf{S}$. The idea of this paragraph is to reformulate the operator A in terms of the sum $\frac{1}{2}(h(\cdot, +\omega_0) + h(\cdot, -\omega_0))$ and the difference $\frac{1}{2}(h(\cdot, +\omega_0) - h(\cdot, -\omega_0))$; namely, we define the following 2×2 invertible matrix:

$$M := \frac{1}{2}\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix},$$

and for $h \in \mathcal{D}$, let $\binom{u}{v} := M \cdot h$, namely

$$\begin{cases} u(\cdot) & := & \frac{1}{2}(h(\cdot,+\omega_0)+h(\cdot,-\omega_0)), \\ v(\cdot) & := & \frac{1}{2}(h(\cdot,+\omega_0)-h(\cdot,-\omega_0)). \end{cases} \tag{4.3}$$

We are now able to define the following operator: $\tilde{A} := M \circ A \circ M^{-1}$, defined on the domain $\widetilde{\mathcal{D}}$

$$\widetilde{\mathcal{D}} := \left\{ (u,v) \in \mathcal{C}^2(\mathbf{S}) \times \mathcal{C}^2(\mathbf{S});\ \int_{\mathbf{S}} u(\theta)\,\mathrm{d}\theta = 0 \right\}, \tag{4.4}$$

given by

$$\forall (u,v) \in \widetilde{\mathcal{D}}, \quad \tilde{A}\begin{pmatrix} u \\ v \end{pmatrix} := \begin{pmatrix} \tilde{A}_1 u \\ \tilde{A}_2 v \end{pmatrix} := \begin{pmatrix} \frac{1}{2}\partial_{\theta^2} u - \partial_\theta\left[u(J * q_0) + q_0(J * u)\right] \\ \frac{1}{2}\partial_{\theta^2} v - \partial_\theta\left[v(J * q_0)\right] \end{pmatrix}. \tag{4.5}$$

The remarkable observation is that operator $\tilde{A}$ is now *uncoupled* w.r.t. variables u and v; consequently, in order to diagonalize $\tilde{A}$, it suffices to diagonalize both components of $\tilde{A}$, namely $\tilde{A}_1$ and $\tilde{A}_2$. This is the purpose of Propositions 4.1 and 4.2.

We make use here of the weighted Sobolev norms $\|\cdot\|_{-1,k}$ defined in (2.7) for different choices of $k(\cdot)$: an important remark is that the first component $\tilde{A}_1$ (with domain $\{u \in \mathcal{C}^2(\mathbf{S}),\ \int_{\mathbf{S}} u = 0\}$) is exactly equal to the McKean-Vlasov operator L_{q_0} with no disorder defined in (2.11). Following § 2.3 and [5], the natural space in where to study the operator $\tilde{A}_1$ is the space $\mathbf{H}^{-1}_{q_0}$ defined in (2.7), for the weight $k(\cdot) = q_0(\cdot)$ the non-disordered stationary solution (recall (1.18)). In this space, we have

Proposition 4.1. *In $\mathbf{H}^{-1}_{q_0}$, $\tilde{A}_1$ is essentially self-adjoint with compact resolvent and spectrum in the negative part of the real axis. 0 is a one-dimensional eigenvalue, spanned by $\partial_\theta q_0$. The spectral gap $\lambda_K(\tilde{A}_1) = \lambda_K(L_{q_0})$ between 0 and the rest of the spectrum is strictly positive.*

Moreover, the self-adjoint extension of $\tilde{A}_1$ is the infinitesimal generator of a strongly continuous semi-group of contractions $\tilde{T}_1(t)$ on $\mathbf{H}^{-1}_{q_0}$. For every $0 < \alpha < \frac{\pi}{2}$, this semigroup can be extended to an analytic semigroup $\tilde{T}_1(z)$ defined on $\Delta_\alpha = \{\lambda;\ |\arg(\lambda)| < \alpha\}$ and one has the following estimate on its resolvent (where $\Sigma_\alpha = \left\{\lambda \in \mathbf{C};\ |\arg(\lambda)| < \frac{\pi}{2} + \alpha\right\} \cup \{0\}$):

$$\forall \alpha \in \left(0, \frac{\pi}{2}\right),\ \forall \lambda \in \Sigma_\alpha,\ \left\| R(\lambda, \tilde{A}_1) \right\|_{-1,q_0} \leqslant \frac{1}{1-\sin(\alpha)} \cdot \frac{1}{|\lambda|}. \tag{4.6}$$

The second component $\tilde{A}_2$ is a second order ordinary differential operator, with domain $\mathcal{C}^2(\mathbf{S})$. As we will see in § 4.1.2, the natural space in which to study $\tilde{A}_2$ is the space $\mathbf{H}^{-1}_w$, for the choice of the weight function $\theta \mapsto w(\theta) = \frac{e^{-\Phi(\theta)}}{\int_{\mathbf{S}} e^{-\Phi}}$, with

$$\Phi(\theta) := -2Kr_0\cos(\theta), \tag{4.7}$$

where r_0 is given by (1.19). Namely, we have

Proposition 4.2. *The operator $(\tilde{A}_2, \mathcal{C}^2(\mathbf{S}))$ is essentially self-adjoint in $\mathbf{H}^{-1}_w$ and has compact resolvent. Hence, its spectrum consists of isolated eigenvalues with finite multiplicities. The kernel of $\tilde{A}_2$ is of dimension 1, spanned by $w(\theta) = \frac{e^{-\Phi(\theta)}}{\int_{\mathbf{S}} e^{-\Phi}}$. Moreover, we have the following spectral gap estimation:*

$$\forall v \in \mathcal{C}^2(\mathbf{S}), \quad -\left\langle \tilde{A}_2 v\,,\, v \right\rangle_{-1,w} \geqslant \frac{e^{-4Kr_0}}{2} \left\| v - \left(\int_{\mathbf{S}} v\right) w \right\|_{-1,w}, \tag{4.8}$$

so that the spectrum of $\tilde{A}_2$ *lies in the negative part of the real axis and the distance between* 0 *and the rest of the spectrum* $\lambda_K(\tilde{A}_2)$ *is at least equal to* $\frac{e^{-4Kr_0}}{2}$. *One also has explicit estimate on the resolvent of* $\tilde{A}_2$*:*

$$\forall \alpha \in (0, \frac{\pi}{2}),\ \forall \lambda \in \Sigma_\alpha,\ \left\| R(\lambda, \tilde{A}_2) \right\|_{-1,w} \leqslant \frac{1}{1 - \sin(\alpha)} \cdot \frac{1}{|\lambda|}. \tag{4.9}$$

Putting things together, the natural norm for the operator $\tilde{A} = (\tilde{A}_1, \tilde{A}_2)$ is the Hilbert-norm, for $(u, v) \in \widetilde{\mathcal{D}}$: $\left(\| u \|_{-1,q_0}^2 + \| v \|_{-1,w}^2\right)^{\frac{1}{2}}$. But since $\tilde{A}$ is the conjugate of A through the invertible matrix M, to say that $\tilde{A}$ is essentially self-adjoint for the previous norm is equivalent to say that A is essentially self-adjoint for the corresponding conjugate norm:

$$\forall h \in \mathcal{D},\ \| h \|_{H_w} := \left(\left\| \frac{1}{2} \left(h(\cdot, +\omega_0) + h(\cdot, -\omega_0) \right) \right\|_{-1,q_0}^2 + \left\| \frac{1}{2} \left(h(\cdot, +\omega_0) - h(\cdot, -\omega_0) \right) \right\|_{-1,w}^2 \right)^{\frac{1}{2}}. \tag{4.10}$$

The results of § 4.1 can be summed-up in the following proposition, which is an easy consequence of Propositions 4.1 and 4.2:

Proposition 4.3. *For the norm* $\| \cdot \|_{H_w}$ *defined in* (4.10)*, the operator* $(A, \mathcal{D})$ *is essentially self-adjoint, with compact resolvent. In particular, the spectrum of (the self-adjoint extension of)* A *is pure-point, and consists of eigenvalues with finite multiplicities. Moreover it lies in the negative part of the real-axis and* A *is the infinitesimal generator of an analytic semigroup of operators* $T_A(z)$ *defined on a domain* $\Delta_\alpha = \{z \in \mathbf{C}; |\arg(z)| < \alpha\}$*, for any* $0 < \alpha < \frac{\pi}{2}$. *One also has the following estimate about the resolvent of* A*:*

$$\forall \alpha \in \left(0, \frac{\pi}{2}\right),\ \forall \lambda \in \Sigma_\alpha,\ \| R(\lambda, A) \|_{H_w} \leqslant \frac{1}{1 - \sin(\alpha)} \cdot \frac{1}{|\lambda|}. \tag{4.11}$$

The kernel of A *is of dimension* 2*, spanned by* $\left\{\partial_\theta q_0 + \frac{e^{-\Phi}}{\int_{\mathbf{S}} e^{-\Phi}}, \partial_\theta q_0 - \frac{e^{-\Phi}}{\int_{\mathbf{S}} e^{-\Phi}}\right\}$ *and the eigenvalue* 0 *is separated from the rest of the spectrum with a distance* $\lambda_K(A) := \min\left(\lambda_K(\tilde{A}_1), \lambda_K(\tilde{A}_2)\right)$*, where* $\lambda_K(\tilde{A}_1)$ *and* $\lambda_K(\tilde{A}_2)$ *are defined in Propositions 4.1 and 4.2 respectively.*

Remark 4.4. *Note that this norm* $\| \cdot \|_{H_w}$ *is equivalent to the norm* $\| \cdot \|_{\mathbf{H}_\mu}$ *defined in § 2.2.2, since the weights* q_0 *and* w *are bounded above and below. In particular, in the space* $\mathbf{H}_\mu^{-1}$*, the operator* A *(although no longer self-adjoint) still generates an analytic semi-group with the same spectrum and the same spectral gap.*

The aim of paragraphs § 4.1.1 (resp. § 4.1.2) is to prove Proposition 4.1 (resp. Proposition 4.2).

4.1.1. *Spectral properties of* $\tilde{A}_1$*: proof of Proposition 4.1.* Much about the operator $\tilde{A}_1$ has already been done since it exactly corresponds to the linear evolution operator L_{q_0} of the non-disordered Kuramoto model studied in [5]: we know from Proposition 2.4 that $\tilde{A}_1$ is essentially self-adjoint and dissipative in $\mathbf{H}_{q_0}^{-1}$. It remains to prove that $\tilde{A}_1$ generates an analytic semigroup $\tilde{T}_1(t)$ in an appropriate domain. We refer to classical references [20, 27, 33] for detailed definitions of analytic semigroups of operators defined on a sector of the complex plane. In particular, we recall the result we use concerning analytic extensions of strongly continuous semigroups. Its proof can be found in [33, Th 5.2, p.61].

Proposition 4.5. *Let $T(t)$ a uniformly bounded strongly continuous semigroup, whose infinitesimal generator F is such that $0 \in \rho(F)$ and let $\alpha \in (0, \frac{\pi}{2})$. The following statements are equivalent:*

1. *$T(t)$ can be extended to an analytic semigroup in the sector $\Delta_\alpha = \{\lambda \in \mathbf{C}; |\arg(\lambda)| < \alpha\}$ and $\|T(z)\|$ is uniformly bounded in every closed sub-sector $\bar{\Delta}_{\alpha'}$, $\alpha' < \alpha$, of Δ_α,*
2. *There exists $M > 0$ such that*

$$\rho(F) \supset \Sigma_\alpha = \left\{\lambda \in \mathbf{C}; |\arg(\lambda)| < \frac{\pi}{2} + \alpha\right\} \cup \{0\}, \tag{4.12}$$

and

$$\|R(\lambda, F)\| \leqslant \frac{M}{|\lambda|}, \quad \lambda \in \Sigma, \lambda \neq 0\,. \tag{4.13}$$

We are now in position to prove the rest of Proposition 4.1: we know from [5, Prop. 2.3, Prop. 2.6] that for any $\lambda > 0$, $\lambda - \tilde{A}_1$ is positive with range $\mathbf{H}^{-1}_{q_0}$. Consequently, we are in position to apply Lumer-Phillips Theorem (see [33, Th 4.3 p.14]): $\tilde{A}_1$ is the infinitesimal generator of a C_0 semi-group of contractions denoted by $\tilde{T}_1(t)$.

The rest of the proof is devoted to show the existence of an analytic extension of this semigroup in a proper sector. We follow here the lines of the proof of [33, Th 5.2, p. 61-62], but with explicit estimates on the resolvent: let us first replace the operator $\tilde{A}_1$ by a small perturbation: for all $\varepsilon > 0$, let $\tilde{A}_{1,\varepsilon} := \tilde{A}_1 - \varepsilon$, so that 0 belongs to $\rho(\tilde{A}_{1,\varepsilon})$. The operator $\tilde{A}_{1,\varepsilon}$ has the following properties: as $\tilde{A}_1$, it is self-adjoint and generates a strongly continuous semigroup of operators (which is $\tilde{T}_{1,\varepsilon}(t) = \tilde{T}_1(t)e^{-\varepsilon t}$). Moreover, since $\tilde{A}_{1,\varepsilon}$ is self-adjoint, we have

$$\forall \lambda \in \mathbf{C} \smallsetminus \mathbf{R}, \left\| R(\lambda, \tilde{A}_{1,\varepsilon}) \right\|_{-1,q_0} \leqslant \frac{1}{|\Im(\lambda)|}, \tag{4.14}$$

and since the spectrum of $\tilde{A}_{1,\varepsilon}$ is negative, for every $\lambda \in \mathbf{C}$ such that $\Re(\lambda) > 0$, we have

$$\left\| R(\lambda, \tilde{A}_{1,\varepsilon}) \right\|_{-1,q_0} \leqslant \frac{1}{|\lambda|}. \tag{4.15}$$

Let us prove that for $\lambda \in \Sigma_\alpha$,

$$\left\| R(\lambda, \tilde{A}_{1,\varepsilon}) \right\|_{-1,q_0} \leqslant \frac{1}{1 - \sin(\alpha)} \cdot \frac{1}{|\lambda|}\,. \tag{4.16}$$

Note that (4.16) is clear from (4.14) and (4.15) when λ is such that $\Re(\lambda) \geqslant 0$. Let us prove it for $\lambda \in \Sigma_\alpha$ with $\Re(\lambda) < 0$. Consider $\sigma > 0, \tau \in \mathbb{R}$ to be chosen appropriately later and write the following Taylor expansion for $R(\lambda, \tilde{A}_\varepsilon)$ around $\sigma + i\tau$ (at least well defined in a neighborhood of $\sigma + i\tau$ since $\sigma > 0$):

$$R(\lambda, \tilde{A}_{1,\varepsilon}) = \sum_{n=0}^{\infty} R(\sigma + i\tau, \tilde{A}_{1,\varepsilon})^{n+1} ((\sigma + i\tau) - \lambda)^n\,. \tag{4.17}$$

This series $R(\cdot, \tilde{A}_{1,\varepsilon})$ is well defined in $\lambda \in \Sigma_\alpha$ with $\Re(\lambda) < 0$ if one can choose σ, τ and $k \in (0,1)$ such that $\left\| R(\sigma + i\tau, \tilde{A}_{1,\varepsilon}) \right\|_{-1,q_0} |\lambda - (\sigma + i\tau)| \leqslant k < 1$. In particular, using (4.14), it suffices to have $|\lambda - (\sigma + i\tau)| \leqslant k|\tau|$ and since $\sigma > 0$ is arbitrary, it suffices to find $k \in (0,1)$ and τ with $|\lambda - i\tau| \leqslant k|\tau|$ to obtain the convergence of (4.17). For this $\lambda \in \Sigma_\alpha$ with $\Re(\lambda) < 0$, let us define λ' and τ as in Figure 4. Then, $|\lambda - i\tau| \leqslant |\lambda' - i\tau| = \sin(\alpha)|\tau|$ with $\sin(\alpha) \in (0,1)$. So the series converges for $\lambda \in \Sigma_\alpha$ and one has, using again (4.14),

$$\left\| R(\lambda, \tilde{A}_{1,\varepsilon}) \right\|_{-1,q_0} \leqslant \frac{1}{(1-\sin(\alpha))|\tau|} \leqslant \frac{1}{1-\sin(\alpha)} \cdot \frac{1}{|\lambda|}. \tag{4.18}$$

The fact that $\tilde{T}_{1,\varepsilon}(t)$ can be extended to an analytic semigroup $\tilde{T}_{1,\varepsilon}(z)$ on the domain Δ_α is a simple application of (4.16) and Proposition 4.5, with $M := \frac{1}{1-\sin(\alpha)}$. Let us then define $\tilde{T}_1(z) := e^{\varepsilon z}\tilde{T}_{1,\varepsilon}(z)$, for $z \in \Delta_\alpha$ so that $\tilde{T}_1(z)$ is an analytic extension of $\tilde{T}_1(t)$ (an argument of analyticity shows that $\tilde{T}_1(z)$ does not depend on ε).



FIGURE 4. The set Σ_α.

Note that estimation (4.6) can be obtained by letting $\varepsilon \to 0$ in (4.16). □

4.1.2. *Spectral properties of $\tilde{A}_2$: proof of Proposition 4.2.* $\tilde{A}_2$ may be written as

$$\tilde{A}_2 v = \frac{1}{2}\partial_\theta^2 v + \partial_\theta \left(v K r_0 \sin(\cdot) \right), \tag{4.19}$$

where $r_0 = \Psi_0(2Kr_0)$ (recall (1.19)).

One recognizes in $\tilde{A}_2$ a second order differential operator on $\mathcal{C}^2(\mathbf{S})$ which can be identified to a simple instance of a Fokker-Planck operator with a sine potential (see [35]). This operator can easily be seen, by integrations by parts in an appropriate weighted $\mathbf{L}^2$-space as a Sturm-Liouville operator acting on $\mathcal{C}^2$, 2π-periodic functions. A vast literature exists on the spectral properties of such operators (see [30, 12, 8, 31, 3, 14]). But the problem is that a $\mathbf{L}^2$-norm is not appropriate for the future study of the SPDE (1.7): a look at the covariance structure of the noise W (see (1.9)) shows that W naturally lives in a $\mathbf{H}^{-1}$-space instead of a $\mathbf{L}^2$-space.

An easy calculation shows that $\tilde{A}_2$ can be rewritten in terms of the weight function Φ defined in (4.7):

$$\tilde{A}_2 v = \frac{1}{2}\partial_\theta \left(e^{-\Phi} \partial_\theta \left(e^{\Phi} v \right) \right). \tag{4.20}$$

Let w be:

$$w(\theta) := \frac{e^{-\Phi(\theta)}}{\int_{\mathbf{S}} e^{-\Phi}}. \tag{4.21}$$

One directly sees from (4.20) that w lies in the kernel of $\tilde{A}_2$: $\tilde{A}_2 w = 0$.

We are now in position to prove Proposition 4.2: let us place ourselves in the framework of the weighted-Sobolev spaces $\left(\mathbf{L}^2_w, \langle \cdot\,,\,\cdot \rangle_{2,w}\right)$ and $\left(\mathbf{H}^{-1}_w, \langle \cdot\,,\,\cdot \rangle_{-1,w}\right)$ constructed in § 2.2.1, in the particular case of $k(\cdot) = w(\cdot)$.

Proof of Proposition 4.2. In $\mathbf{H}^{-1}_w$, the operator $(\tilde{A}_2, \mathcal{C}^2(\mathbf{S}))$ is formally symmetric: indeed, for u, $v \in \mathcal{C}^2(\mathbf{S})$, for u_0 ad v_0 defined by (2.6), we have successively,

$$\begin{aligned}\left\langle \tilde{A}_2 u\,,\, v\right\rangle_{-1,w} &= \left(\int_{\mathbf{S}} e^{-\Phi}\right)\int_{\mathbf{S}} e^{\Phi}\left(\frac{1}{2}e^{-\Phi}\partial_\theta\left(e^{\Phi}u\right)\right)\mathcal{V}_0 = -\frac{1}{2}\left(\int_{\mathbf{S}} e^{-\Phi}\right)\int_{\mathbf{S}} e^{\Phi} u v_0,\\ &= -\frac{1}{2}\int_{\mathbf{S}}\frac{u_0 v_0}{w} - \frac{1}{2}\int_{\mathbf{S}} u\int_{\mathbf{S}} v_0 = -\frac{1}{2}\int_{\mathbf{S}}\frac{u_0 v_0}{w}.\end{aligned} \tag{4.22}$$

The latter expression being symmetric in u and v.

Let us prove that $(\tilde{A}_2, \mathcal{C}^2(\mathbf{S}))$ is essentially self-adjoint: let $\mathcal{E}_2$ be the following Dirichlet form

$$\mathcal{E}_2(u,v) := \left\langle u\,,\,(1-\tilde{A}_2)v\right\rangle_{-1,w} = \int_{\mathbf{S}} u\int_{\mathbf{S}} v + \int_{\mathbf{S}}\frac{\mathcal{V}_0\mathcal{U}_0}{w} + \frac{1}{2}\int_{\mathbf{S}}\frac{u_0 v_0}{w}. \tag{4.23}$$

Then it is easy to see that $\mathcal{E}_2$ is a continuous bilinear form on $\mathbf{L}^2_w$ (thanks to Poincaré inequality). Moreover $\mathcal{E}_2$ is coercive, namely, for all $u \in \mathbf{L}^2_w$:

$$\begin{aligned}\mathcal{E}_2(u,u) &= \left(\int_{\mathbf{S}} u\right)^2 + \int_{\mathbf{S}}\frac{\mathcal{U}_0^2}{w} + \frac{1}{2}\int_{\mathbf{S}}\frac{u_0^2}{w},\\ &\geqslant \left(\int_{\mathbf{S}} u\right)^2 + \frac{1}{2}\left\| u - \left(\int_{\mathbf{S}} u\right) w\right\|^2_{2,w} \geqslant \frac{1}{2}\left\| u\right\|^2_{2,w}.\end{aligned} \tag{4.24}$$

Since for all $f \in \mathbf{H}^{-1}_w$, the linear form $v \mapsto \langle v\,,\, f\rangle_{-1,w}$ is continuous on $\mathbf{L}^2_w$, an application of Lax-Milgram Theorem shows that for such an $f \in \mathbf{H}^{-1}_w$ there exists an unique $u \in \mathbf{L}^2_w$ such that for all $v \in \mathbf{L}^2_w$

$$\mathcal{E}_2(v,u) = \langle v\,,\, f\rangle_{-1,w}. \tag{4.25}$$

It is then easy to see that $\int_{\mathbf{S}} f = \int_{\mathbf{S}} u$ and that for almost every $\theta \in \mathbf{S}$,

$$\frac{1}{2}\frac{u_0(\theta)}{w(\theta)} = -\int_0^\theta \frac{\mathcal{F}_0}{w} + \int_0^\theta \frac{\mathcal{U}_0}{w}. \tag{4.26}$$

Since $u \in \mathbf{L}^2_w$, $\mathcal{U}_0$ admits a $\mathcal{C}^1$-version and if we assume that f is square-integrable, the same argument holds for the first term of the right-hand side of (4.26). So, if f is square integrable, u_0 admits a $\mathcal{C}^2$-version. To sum-up, if we suppose that f is continuous, there exists $u \in \mathcal{C}^2(\mathbf{S})$ such that, applying $\partial_\theta\left(e^{-\Phi}\partial_\theta(\cdot)\right)$ to (4.26):

$$\begin{aligned}f = f_0 + \left(\int_{\mathbf{S}} f\right) w &= -\frac{1}{2}\partial_\theta\left(e^{-\Phi}\partial_\theta\left(e^{\Phi}u_0\right)\right) + u_0 + p_w(u)w,\\ &= -\tilde{A}_2 u_0 + u = (1-\tilde{A}_2)u.\end{aligned} \tag{4.27}$$

But since those functions f are dense in $\mathbf{H}^{-1}_w$, we see that the range of $1-\tilde{A}_2$ is dense so that $\tilde{A}_2$ is essentially self-adjoint.

Secondly, the spectral gap estimation (4.8) holds: indeed, for every $u \in \mathcal{C}^2(\mathbf{S})$, we have using (4.22) and Poincaré inequality:

$$\begin{aligned} -\left\langle \tilde{A}_2 v\,,\, v\right\rangle_{-1,w} &= \frac{1}{2}\left(\int_{\mathbf{S}} e^{-\Phi}\right)\int_{\mathbf{S}} e^{\Phi} v_0^2,\\ &\geqslant \frac{1}{2}e^{-2Kr_0}\left(\int_{\mathbf{S}} e^{-\Phi}\right)\int_{\mathbf{S}} \mathcal{V}_0^2 \geqslant \frac{1}{2}e^{-4Kr_0}\left(\int_{\mathbf{S}} e^{-\Phi}\right)\int_{\mathbf{S}} e^{\Phi}\mathcal{V}_0^2,\\ &= \frac{1}{2}e^{-4Kr_0}\left\| v - \left(\int_{\mathbf{S}} v\right) w\right\|_{-1,w}^2 . \end{aligned}$$

Moreover, $\tilde{A}_2$ has compact resolvent: it suffices to prove that $\lambda - \tilde{A}_2$ has compact resolvent for at least one value of λ. We prove it for $\lambda = 1$ which is indeed in the resolvent set, thanks to the beginning of this proof. For $u \in \mathbf{H}_w^{-1}$, let us consider $f := (1-\tilde{A}_2)^{-1}u$ so that $\left\langle f\,,\,(1-\tilde{A}_2)f\right\rangle_{-1,w} = \langle f\,,\,u\rangle_{-1,w}$. Using the coerciveness of $\mathcal{E}_2$, one has for some constant c, $c\,\|\, f\,\|_{2,w}^2 \leqslant \langle f\,,\,u\rangle_{-1,w} \leqslant \|\, f\,\|_{-1,w}\,\|\, u\,\|_{-1,w}$. Using the continuous injection of $\mathbf{L}_w^2$ into $\mathbf{H}_w^{-1}$ (say $\|\cdot\|_{-1,w} \leqslant C\,\|\cdot\|_{2,w}$, for some positive constant C), one has

$$\|\, f\,\|_{2,w} \leqslant \frac{C}{c}\,\|\, u\,\|_{-1,w} . \tag{4.28}$$

So $(1-\tilde{A}_2)^{-1}$ maps sequences that are bounded in $\mathbf{H}_w^{-1}$ into sequences that are bounded in $\mathbf{L}_w^2$. It remains then to prove that the injection of $\mathbf{L}_w^2$ into $\mathbf{H}_w^{-1}$ is compact. This is indeed true since for every $v \in \mathbf{H}_w^{-1}$, one has, by Cauchy-Schwartz inequality $|\mathcal{V}_0(\theta) - \mathcal{V}_0(\theta')| \leqslant C\,\|\, v_0\,\|_{2,w}\sqrt{|\theta-\theta'|} \leqslant C\,\|\, v\,\|_{2,w}\sqrt{|\theta-\theta'|}$. That means that, by Ascoli-Arzela Theorem that the sets $\left\{v \in \mathbf{H}_w^{-1};\ \|\, v\,\|_{2,w} \leqslant cst\right\}$ are relatively compact in $\mathcal{C}(\mathbf{S})$ and also in $\mathbf{L}_w^2$. That completes the proof.

The fact that $\tilde{A}_2$ generates an analytic semigroup $\tilde{T}_2(z)$ on the same sector Δ_α as well as estimation (4.9) can be derived in the same way as in § 4.1.1. That concludes the proof of Proposition 4.2. □

4.2. **Control on the perturbation B.** In order to derive spectral properties for the operator $L = A + B$, we need to have a precise estimation about the smallness of the perturbation B w.r.t. operator A studied in the previous paragraph § 4.1.

Remark 4.6. Note that for simplicity, we work now with the norm $\|\cdot\|_{\mathbf{H}_\mu}$ (recall Remark 2.3); as already mentioned this norm is equivalent to the norm $\|\cdot\|_{H_w}$ used in § 4.1. Recall also the definition of the space $(\mathbf{L}^2, \|\cdot\|_2)$ defined in Remark 2.2 and of $(\mathbf{L}_\mu^2, \|\cdot\|_{\mu,2})$ defined in Remark 2.3.

Secondly, since the whole operator L is no longer symmetric in $\mathbf{H}_\mu^{-1}$, its spectrum need not be real. Thus, we will assume for the rest of this document that we work with the complexified versions of the scalar products defined previously in this paper. The results concerning the operator A are obviously still valid.

The smallness of the perturbation B with respect to A can be quantified in terms of the difference $\|\, q(\cdot,\omega) - q_0(\cdot)\,\|_\infty$, $\omega \in \mathrm{Supp}(\mu)$. For the ease of exposition, we do not attempt to derive precise estimations of this difference $\|\, q(\cdot,\omega) - q_0(\cdot)\,\|_\infty$ (Lemma 4.7) and of coefficients $a(\omega_0)$ and $b(\omega_0)$ (Lemma 4.8), in terms of the coupling strength K. We will denote as c a positive constant (depending on K) which may change from a line to another.

Lemma 4.7. *For $\omega > 0$ and $K > 1$, let us define*

$$\| q - q_0 \|_\infty := \sup_{\theta \in \mathbf{S}, |u| \leqslant \omega} |q(\theta, u) - q_0(\theta)|. \tag{4.29}$$

Then $\| q - q_0 \|_\infty = O(\omega)$, as $\omega \to 0$.

Proof. This is clear since one can bound $\partial_\omega q(\theta, \omega)$ uniformly in (θ, ω), as $\omega \to 0$ (by a constant depending on K). □

The purpose of the following proposition is to prove the boundedness w.r.t. A of the perturbation B in terms of the size of the disorder:

Proposition 4.8. *The operator B is A-bounded in the sense that there exist positive constants $a(\omega_0) = a(\omega_0, K)$ and $b(\omega_0) = b(\omega_0, K)$ such that*

$$\forall h \in \mathcal{D}, \quad \| Bh \|_{\mathbf{H}_\mu} \leqslant a(\omega_0) \| h \|_{\mathbf{H}_\mu} + b(\omega_0) \| Ah \|_{\mathbf{H}_\mu}, \tag{4.30}$$

and moreover, for fixed $K > 1$,

$$a(\omega_0) = O(\omega_0), \text{ and } b(\omega_0) = O(\omega_0), \quad \text{as } \omega_0 \to 0, \tag{4.31}$$

Proof of Proposition 4.8. Recall that in what follows, $\langle h \rangle_\mu(\cdot) = \int_{\mathbf{R}} h(\cdot, \omega) \mu(\mathrm{d}\omega)$ is the averaging of $h(\cdot, \omega)$ and that c is a constant (depending on K) that may change from a line to another. The proof consists in two steps: let us prove firstly that there exist some constant α_{K,ω_0} such that for all $h \in \mathcal{D}$,

$$\| Bh \|_{\mathbf{H}_\mu} \leqslant \alpha_{K,\omega_0} \| h \|_{\mu,2}, \tag{4.32}$$

Indeed, for given $h \in \mathcal{D}$, for all $\omega \in \mathrm{Supp}(\mu)$, we have $\|Bh(\cdot,\omega)\|_{-1} = \| \mathcal{B}h(\cdot,\omega) \|_2$, where $\mathcal{B}h(\cdot,\omega)$ is the appropriate primitive of $Bh(\cdot,\omega)$ in $\mathbf{H}^{-1}$, (recall Remark 2.2) namely:

$$\begin{aligned} \mathcal{B}h := & -h\left(\langle J * (q - q_0) \rangle_\mu + \omega\right) - (q - q_0) \cdot \langle J * h \rangle_\mu \\ & + \int_{\mathbf{S}} h \cdot \left(\langle J * (q - q_0) \rangle_\mu + \omega\right) + \int_{\mathbf{S}} (q - q_0) \cdot \langle J * h \rangle_\mu. \end{aligned}$$

Using the boundedness of q_0 and the bounds $| (J * \varepsilon) | \leqslant 4K \| \varepsilon \|_\infty$ and $|\langle J * h \rangle_\mu| \leqslant \frac{K}{\sqrt{2}} \| \langle h \rangle_\mu \|_2$, it is easy to deduce that, for some constant $c > 0$:

$$|\mathcal{B}h| \leqslant c(\| q - q_0 \|_\infty + \omega_0)(|h| + \| \langle h \rangle_\mu \|_2). \tag{4.33}$$

Consequently,

$$\| Bh \|_{\mathbf{H}_\mu} = \left(\langle \| \mathcal{B}h \|_2^2 \rangle_\mu\right)^{\frac{1}{2}} \leqslant c(\| q - q_0 \|_\infty + \omega_0) \| h \|_{\mu,2}, \tag{4.34}$$

so that (4.32) is satisfied for some coefficient α_{K,ω_0} such that (Lemma 4.7) $\alpha_{K,\omega_0} = O(\omega_0)$ as $\omega_0 \to 0$.

The second step of the proof is to control the $\mathbf{L}^2$-norm $\| h \|_{\mu,2}$ of h with the $\mathbf{H}_\mu^{-1}$-norms of Ah and h: namely we prove that there exist constants γ_K and δ_K such that

$$\| h \|_{\mu,2} \leqslant \gamma_K \| Ah \|_{\mathbf{H}_\mu} + \delta_K \| h \|_{\mathbf{H}_\mu}. \tag{4.35}$$

The proof is based on a usual interpolation argument: for all integer $n > 1$, for any $f \in \mathcal{C}^2(\mathbf{S})$, one has

$$\| \partial_\theta f \|_2^2 \leqslant \sqrt{n} \| f \|_2 \frac{\| \partial_\theta^2 f \|_2}{\sqrt{n}} \leqslant \frac{n}{2} \| f \|_2^2 + \frac{\| \partial_\theta^2 f \|_2^2}{2n}. \tag{4.36}$$

Let us use this interpolation relation (4.36) to derive (4.35): for all $h \in \mathcal{D}$, $\omega \in \mathrm{Supp}(\mu)$, one has

$$\| h(\cdot,\omega) \|_2^2 = \left(\int_{\mathbf{S}} h(\cdot,\omega) \right)^2 + \| h_0(\cdot,\omega) \|_2^2 \tag{4.37}$$

Applying relation (4.36) with $f(\cdot) = \mathcal{H}_0(\cdot,\omega)$ we obtain

$$\| h(\cdot,\omega) \|_2^2 \leqslant \left| \int_{\mathbf{S}} h(\cdot,\omega) \right|^2 + \frac{n}{2} \| \mathcal{H}_0(\cdot,\omega) \|_2^2 + \frac{\| \partial_\theta h(\cdot,\omega) \|_2^2}{2n}, \tag{4.38}$$

where we used the fact that $\partial_\theta h_0(\cdot,\omega) = \partial_\theta h(\cdot,\omega)$. Integrating w.r.t. μ,

$$\| h \|_{\mu,2}^2 \leqslant \int_{\mathbf{R}} \left| \int_{\mathbf{S}} h \right|^2 \mathrm{d}\mu + \frac{n}{2} \| \mathcal{H}_0 \|_{\mu,2}^2 + \frac{\| \partial_\theta h \|_{\mu,2}^2}{2n}. \tag{4.39}$$

But, as previously for the operator B, a simple calculation shows that for all $\omega \in \mathrm{Supp}(\mu)$, we have $\|Ah(\cdot,\omega)\|_{-1} = \| \mathcal{A}h(\cdot,\omega) \|_2$, where $\mathcal{A}h$ is the appropriate primitive of Ah in $\mathbf{H}^{-1}$ (recall (4.1)):

$$\mathcal{A}h = \frac{1}{2}\partial_\theta h - h(J * q_0) - q_0 \langle J * h \rangle_\mu + \int_{\mathbf{S}} \left(h(J * q_0) + q_0 \langle J * h \rangle_\mu \right), \tag{4.40}$$

so that, for some constant $c > 0$

$$\| \partial_\theta h \|_{\mu,2}^2 \leqslant 12 \| \mathcal{A}h \|_{\mu,2}^2 + c \| h \|_{\mu,2}^2 . \tag{4.41}$$

Injecting this inequality in (4.39), one obtains, for some constant $c > 0$

$$\| h \|_{\mu,2}^2 \leqslant \int_{\mathbf{R}} \left| \int_{\mathbf{S}} h \right|^2 \mathrm{d}\mu + \frac{n}{2} \| \mathcal{H}_0 \|_{\mu,2}^2 + \frac{1}{2n} \left(12 \| \mathcal{A}h \|_{\mu,2}^2 + c \| h \|_{\mu,2}^2 \right). \tag{4.42}$$

Choosing $n > 1$ sufficiently large so that the coefficient in front of $\| h \|_{\mu,2}^2$ in the right-hand side of (4.42) is lower than $\frac{1}{2}$ leads to (for some constant $c > 0$):

$$\| h \|_{\mu,2}^2 \leqslant 2 \int_{\mathbf{R}} \left| \int_{\mathbf{S}} h \right|^2 \mathrm{d}\mu + c \| \mathcal{H}_0 \|_{\mu,2}^2 + c \| \mathcal{A}h \|_{\mu,2}^2 \leqslant c \| h \|_{\mathbf{H}_\mu}^2 + c \| Ah \|_{\mathbf{H}_\mu}^2 ,$$

which shows (4.35). Putting together estimates (4.32) and (4.35), we find the A-boundedness of B (4.30) with coefficients $a(\omega_0)$ and $b(\omega_0)$ which satisfy (4.31), thanks to Lemma 4.7. ☐

Proposition 4.9. *The operator B is A-compact, in the sense that for any sequence $(h_p)_{p \geqslant 0} \in \mathcal{D}^{\mathbf{N}}$ such that $\| h_p \|_{\mathbf{H}_\mu}$ and $\| Ah_p \|_{\mathbf{H}_\mu}$ are bounded, there exists a convergent subsequence for $(Bh_p)_{p \geqslant 1}$.*

Proof of Proposition 4.9. Let $(h_p)_{p \geqslant 0}$ a sequence in $\mathcal{D}$ such that $\| h_p \|_{\mathbf{H}_\mu}$ and $\| Ah_p \|_{\mathbf{H}_\mu}$ are bounded by a constant c. A closer look at the operator B defined in (4.2) and the definition of the norm $\| \cdot \|_{\mathbf{H}_\mu}$ in (2.8) shows that it suffices to prove that there exists a subsequence $(h_{p_k})_{k \geqslant 0}$ that is convergent in $\mathbf{L}^2_\mu$. In particular, for all $p \geqslant 0$, $\| Ah_p \|_{\mathbf{H}_\mu} \leqslant c$. Using this boundedness and (4.41), we have $\| \partial_\theta h_p \|_{\mu,2} \leqslant c + c \| h_p \|_{\mu,2}$, so that

$$\| \partial_\theta h_p \|_{\mu,2} \leqslant c + c \left(\int_{\mathbf{R}} \left| \int_{\mathbf{S}} h \right|^2 \mathrm{d}\mu + \| h_{0,p} \|_{\mu,2}^2 \right), \tag{4.43}$$

$$\leqslant c + c \left(\int_{\mathbf{R}} \left| \int_{\mathbf{S}} h \right|^2 \mathrm{d}\mu + \frac{n}{2} \| \mathcal{H}_{0,p} \|_{\mu,2}^2 + \frac{\| \partial_\theta h_p \|_{\mu,2}^2}{2n} \right), \tag{4.44}$$

where we used again (4.36) for $f = \mathcal{H}_{0,p}(\cdot, \omega)$. Choosing a sufficiently large $n > 1$ leads to $\| \partial_\theta h_{0,p} \|_{\mu,2} = \| \partial_\theta h_p \|_2 \leqslant c + c \| h_p \|_{\mathbf{H}_\mu} \leqslant c$ for a constant c independent of $p \geqslant 0$. An easy application of Cauchy-Schwartz inequality leads to $|h_{0,p}(\theta, \omega) - h_{0,p}(\theta', \omega)| \leqslant \| \partial_\theta h_{0,p} \|_{\mu,2} \sqrt{|\theta - \theta'|}$, for all $\omega \in \{\pm\omega_0\}$. Since the functions $(h_{0,p})_{p \geqslant 0}$ are such that $\int_{\mathbf{S}} h_{0,p} = 0$ for all $p \geqslant 0$, Ascoli-Arzela Theorem and the previous bound show the existence of a convergent subsequence (h_{0,p_k}) (for each $\omega \in \mathrm{Supp}(\mu)$) in the space of continuous functions on $\mathbf{S}$. In particular, this subsequence is convergent in $\mathbf{L}^2_\mu$ and is renamed $(h_{0,p})_{p \geqslant 0}$, with a slight abuse of notations.

The fact that $\int_{\mathbf{R}} | \int_{\mathbf{S}} h_p | \mathrm{d}\mu \leqslant c$ shows that one can extract a further subsequence of $(h_p)_{p \geqslant 0}$ which is also convergent in $\mathbf{L}^2_\mu$. This concludes the proof. □

4.3. **Spectral properties of $L = A + B$.**

We are now in position to extent the spectral properties we obtained in § 4.1 for the operator A to the perturbed operator $L = A + B$, relying on the control we have the perturbation B (see § 4.2) and on the theory of perturbation of operators ([23]) and analytic semi-groups ([33]). Recall again that the disorder is supposed to be binary (see (2.4)).

4.3.1. *The spectrum of L is pure point.*

Proposition 4.10. *For all K, for all $\omega_0 > 0$,*

1. *the operator $(L, \mathcal{D})$ is closable. In that case, its closure has the same domain as the closure of A,*
2. *the closure of L has compact resolvent. In particular, its spectrum is pure point.*

Remark 4.11. *Note that Proposition 4.10 is valid without any assumption on the smallness of ω_0, since it relies on the relative compactness of B with respect to A (Prop. 4.9).*

Proof of Proposition 4.10. It is a simple consequence of the relative compactness of B w.r.t. the self-adjoint operator A. The first assertion of Proposition 4.10 is a consequence of [23, Th. 1.11 p.194] and the second assertion can be found in [28, Lemma 3.6, p.17] for example. □

4.3.2. *L generates an analytic operator.*

We prove mainly that the perturbed operator L generates an analytic semigroup of operators on a appropriate sector. An immediate corollary is the position of the spectrum in a cone whose vertex is zero. We know (Proposition 4.3) that for all $0 < \alpha < \frac{\pi}{2}$, A generates a semigroup of operators on $\Delta_\alpha = \{z \in \mathbf{C};\ |\arg(z)| < \alpha\}$.

Proposition 4.12. *For all $K > 1$, $\varepsilon > 0$ and $0 < \alpha < \frac{\pi}{2}$, there exists $\omega_1 > 0$ (depending on α, K and ε) such that for all $0 < \omega_0 < \omega_1$, the spectrum of L lies within the sector $\Theta_{\varepsilon,\alpha} := \left\{\lambda \in \mathbf{C};\ \frac{\pi}{2} + \alpha \leqslant \arg(\lambda) \leqslant \frac{3\pi}{2} - \alpha\right\} \cup \{\lambda \in \mathbf{C};\ |\lambda| \leqslant \varepsilon\}$. For such ω_0, L generates an analytic semigroup of operators on $\Delta_{\alpha'}$, for some $\alpha' \in (0, \frac{\pi}{2})$.*

Proof of Proposition 4.12. Let $0 < \alpha < \frac{\pi}{2}$ be fixed. Thanks to (4.11), there exists a constant $c > 0$ (which comes from the equivalence of the norms $\| \cdot \|_{\mathbf{H}_\mu}$ and $\| \cdot \|_{H_w}$) such that for every $\lambda \in \Sigma_\alpha := \left\{\lambda \in \mathbf{C};\ |\arg(\lambda)| < \frac{\pi}{2} + \alpha\right\}$, we have the estimates:

$$\| R(\lambda, A) \|_{\mathbf{H}_\mu} \leqslant \frac{c}{(1 - \sin(\alpha))|\lambda|} \quad \text{and,} \quad \| AR(\lambda, A) \|_{\mathbf{H}_\mu} \leqslant 1 + \frac{c}{(1 - \sin(\alpha))}.$$

Then for $\lambda \in \Sigma_\alpha$, $h \in \mathcal{D}$:

$$\begin{aligned} \| BR(\lambda, A)h \|_{\mathbf{H}_\mu} &\leqslant a(\omega_0) \| R(\lambda, A)h \|_{\mathbf{H}_\mu} + b(\omega_0) \| AR(\lambda, A)h \|_{\mathbf{H}_\mu}, \\ &\leqslant \left(a(\omega_0) \frac{c}{(1 - \sin(\alpha))|\lambda|} + b(\omega_0) \left(1 + \frac{c}{1 - \sin(\alpha)} \right) \right) \| h \|_{\mathbf{H}_\mu}. \end{aligned}$$

Let us fix $\varepsilon > 0$ and choose ω_1 such that:

$$\max\left(\frac{4a(\omega_1)c}{\varepsilon(1-\sin(\alpha))}, 4b(\omega_1)\left(1+\frac{c}{1-\sin(\alpha)}\right)\right) \leqslant 1. \tag{4.45}$$

For this choice of ω_1, for all $0 < \omega_0 < \omega_1$, for any $\lambda \in \Sigma_\alpha$ such that $|\lambda| \geqslant \varepsilon \geqslant \frac{4a(\omega_1)c}{1-\sin(\alpha)}$, we have $\| BR(\lambda, A)h \|_{\mathbf{H}_\mu} \leqslant \frac{1}{2} \| h \|_{\mathbf{H}_\mu}$, and thus the operator $1 - BR(\lambda, A)$ is invertible with $\| 1 - BR(\lambda, A) \|_{\mathbf{H}_\mu} \leqslant 2$. Since it can easily be shown that

$$R(\lambda, A+B) = R(\lambda, A)\left(1 - BR(\lambda, A)\right)^{-1},$$

one deduces the following estimates about the resolvent of the perturbed operator $L = A + B$:

$$\forall \lambda \in \Sigma_\alpha,\ |\lambda| \geqslant \varepsilon,\ \| R(\lambda, L) \|_{\mathbf{H}_\mu} \leqslant \frac{2}{(1-\sin(\alpha))|\lambda|}. \tag{4.46}$$

The fact that the spectrum of L lies within $\Theta_{\varepsilon,\alpha}$ is a straightforward consequence of (4.46). Secondly, (4.46) entails that L generates an analytic semigroup of operators on an appropriate sector. Indeed, if one denotes by $L_\varepsilon := L - \varepsilon$, one deduces from (4.46) that $0 \in \rho(L_{2\varepsilon})$ and that for all $\lambda \in \mathbf{C}$ with $\Re(\lambda) > 0$ (in particular, $|\lambda| < |\lambda + 2\varepsilon|$)

$$\begin{aligned} \| R(\lambda, L_{2\varepsilon}) \|_{\mathbf{H}_\mu} = \| R(\lambda + 2\varepsilon, L) \|_{\mathbf{H}_\mu} &\leqslant \frac{2}{(1-\sin(\alpha))|\lambda + 2\varepsilon|}, \\ &\leqslant \frac{2}{(1-\sin(\alpha))|\lambda|}. \end{aligned} \tag{4.47}$$

Hence, using the same arguments of Taylor expansion as in the proof of Proposition 4.1 and applying Proposition 4.5, one easily sees that $L_{2\varepsilon}$ generates an analytic semigroup in a (a priori) smaller sector $\Delta_{\alpha'}$, where $\alpha' \in (0, \frac{\pi}{2})$ can be chosen as $\alpha' := \frac{1}{2}\arctan\left(\frac{1-\sin(\alpha)}{2}\right)$. But if $L_{2\varepsilon}$ generates an analytic semigroup, so does L. ☐

4.3.3. 0 *is an eigenvalue with multiplicity* 2. Let us fix $K > 1$, $\alpha \in (0, \frac{\pi}{2})$, $\rho \in (0,1)$ and define $\varepsilon = \rho\lambda_K(A)$. Applying Proposition 4.12, we know that for small ω_0 (depending on K, α, ρ), L generates an analytic semigroup on $\Theta_{\varepsilon,\alpha}$. Let $\Theta^+_{\varepsilon,\alpha} := \{\lambda \in \Theta_{\varepsilon,\alpha};\ \Re(\lambda) \geqslant 0\}$ be the subset of $\Theta_{\varepsilon,\alpha}$ which lies in the positive part of the complex plane (see Figure 5).

The purpose of this paragraph is to show that one can choose a perturbation B small enough so that no non-zero eigenvalue of $A + B$ remains in the small set $\Theta^+_{\varepsilon,\alpha}$.

To do so, we proceed by an argument of local perturbation: we know that the distance $\lambda_K(A) = \min\left(\lambda_K(\tilde{A}_1), \lambda_K(\tilde{A}_2)\right)$ between the eigenvalue 0 and the rest of the spectrum of A is strictly positive. In particular, one can separate 0 from the rest of the spectrum of A by a circle $\mathscr{C}$ centered in 0 with radius $\left(\frac{\rho+1}{2}\right)\lambda_K(A)$. Note that the appropriate choice of ε made at the beginning of this paragraph ensures that the interior of $\mathscr{C}$ contains $\Theta^+_{\varepsilon,\alpha}$ (Figure 5).

The main argument is the following: by construction of $\mathscr{C}$, 0 is the only eigenvalue (with multiplicity 2) of the non-perturbed operator A lying in the interior of $\mathscr{C}$. A principle of continuity of eigenvalues shows that, while adding a small enough perturbation B to A, the interior of $\mathscr{C}$ still contains either one eigenvalue with algebraic multiplicity 2 or two eigenvalues with multiplicity 1; those perturbed eigenvalues remain close but are *a priori* not equal to the initial eigenvalue 0 (see Figure 5).

But we already know that for the perturbed operator $L = A + B$, 0 is always an eigenvalue (since $L\partial_\theta q = 0$ and $Lp = \partial_\theta q$, recall Th. 2.5). Therefore, the algebraic multiplicity of 0 for



(A) Position of the spectrum for the self-adjoint operator A

(B) Possible position of the spectrum for the operator L

FIGURE 5. The domains $\Theta_{\varepsilon,\alpha}$ and $\Theta^+_{\varepsilon,\alpha}$ (in light grey). Note that the two dimensional eigenvalue 0 for the operator A (5a) may split in two single eigenvalues for the perturbed operator L (5b). These eigenvalues are the only ones within the circle $\mathscr{C}$. But since we already know that the eigenspace in 0 of L is of dimension 2, 0 is still a double eigenvalue for L, by uniqueness.

the operator L is *at least* 2. By uniqueness, one can conclude that 0 is the *only* element of the spectrum of L within $\mathscr{C}$, and is an eigenvalue with algebraic multiplicity *exactly* 2. In particular, there is no element of the spectrum in the positive part of the complex plane.

In order to make this argument precise, we need to quantify the appropriate size of the perturbation B, by explicit estimates on the resolvent $R(\lambda, A)$ on the circle $\mathscr{C}$:

Lemma 4.13. *There exists some explicit constant* $c_{\mathscr{C}}(K)$ *only dependent on* K*, such that for all* $\lambda \in \mathscr{C}$*,*

$$\| R(\lambda, A) \|_{\mathbf{H}_\mu} \leqslant c_{\mathscr{C}}(K), \tag{4.48}$$

$$\| AR(\lambda, A) \|_{\mathbf{H}_\mu} \leqslant 1 + \left(\frac{\rho+1}{2}\right) \cdot c_{\mathscr{C}}(K). \tag{4.49}$$

One can choose $c_{\mathscr{C}}(K)$ *as* $c_{\mathscr{C}}(K) = \frac{1}{\lambda_K(A)} \max\left(\frac{2}{\rho+1}, \frac{2}{1-\rho}\right)$.

Proof of Lemma 4.13. Applying the spectral theorem (see [12, Th. 3, p.1192]) to the essentially self-adjoint operator A, there exists a spectral measure E vanishing on the complementary of the spectrum of A such that $A = \int_{\mathbf{R}} \lambda \, \mathrm{d}E(\lambda)$. In that extent, one has for any $\zeta \in \mathscr{C}$

$$R(\zeta, A) = \int_{\mathbf{R}} \frac{\mathrm{d}E(\lambda)}{\lambda - \zeta}. \tag{4.50}$$

In particular, for $\zeta \in \mathscr{C}$

$$\| R(\zeta, A) \|_{\mathbf{H}_\mu} \leqslant \sup_{\lambda \in \sigma(A)} \frac{1}{|\lambda - \zeta|} \leqslant \frac{\max\left(\frac{2}{\rho+1}, \frac{2}{1-\rho}\right)}{\lambda_K(A)}. \tag{4.51}$$

The estimation (4.49) is straightforward. □

We are now in position to apply our argument of local continuity of eigenvalues: following [23, Th III-6.17, p.178], there exists a decomposition of the operator A according to $\mathbf{H}_\mu^{-1} = F_0 \oplus F_{<0}$ (in the sense that $AF_0 \subset F_0$, $AF_{<0} \subset F_{<0}$ and $P_0\mathcal{D} \subset \mathcal{D}$, where P_0 is the projection on F_0 along $F_{<0}$) in such a way that A restricted to F_0 has spectrum $\{0\}$ and A restricted to $F_{<0}$ has spectrum $\sigma(A) \setminus \{0\} \subseteq \{\lambda \in \mathbf{C}; \Re(\lambda) < 0\}$. Let us note that the dimension of F_0 is *exactly* 2, since the characteristic space of A for the eigenvalue 0 is reduced to its kernel which is of dimension 2 (see Prop. 4.3).

Then, applying [23, Th. IV-3.18, p.214], and using Proposition 4.8, we find that if one chooses $\omega_2 > 0$, such that

$$\sup_{\lambda \in \mathscr{C}} \Big(a(\omega_2) \| R(\lambda, A) \|_{\mathbf{H}_\mu} + b(\omega_2) \| AR(\lambda, A) \|_{\mathbf{H}_\mu} \Big) < 1, \tag{4.52}$$

then for all $0 < \omega_0 < \omega_2$, the perturbed operator L is likewise decomposed according to $\mathbf{H}_\mu^{-1} = G_0 \oplus G_{<0}$, in such a way that $\dim(F_0) = \dim(G_0) = 2$, and that the spectrum of L is again separated in two parts by $\mathscr{C}$. But thanks to Theorem 2.5, we already know that the characteristic space of the perturbed operator L according to the eigenvalue 0 is *at least* of dimension 2 (since $L\partial_\theta q = 0$ and $Lp = \partial_\theta q$). We can conclude, that for such an $0 < \omega_0 < \omega_2$, 0 is the only eigenvalue in $\mathscr{C}$ and that $\dim(G_0)$ is *exactly* 2.

Applying Lemma 4.13, we see that condition (4.52) is satisfied if we choose $\omega_2 > 0$ so that:

$$a(\omega_2) c_{\mathscr{C}}(K) + b(\omega_2) \left(1 + \left(\frac{\rho + 1}{2} \right) c_{\mathscr{C}}(K) \right) < 1. \tag{4.53}$$

In particular, in that case, the spectrum of L is contained in $\{z \in \mathbf{C}; \Re(z) \leqslant 0\}$.

Theorem 2.7 is proved.

5. NON SELF-AVERAGING PHENOMENON FOR THE FLUCTUATION PROCESS (PROOF OF THEOREM 2.9)

The purpose of this section is to prove our main result Theorem 2.9, that is the linear asymptotics (2.19) for the SPDE (1.7).

In our framework, (recall that $\mu = \frac{1}{2}(\delta_{-\omega_0} + \delta_{\omega_0})$), the solution η of evolution (1.7) in $\mathcal{S}'(\mathbf{S} \times \mathbf{R})$ acts on test functions φ of the form $\varphi = (\varphi(\cdot, +\omega_0), \varphi(\cdot, -\omega_0))$. In particular, one can understand η as an element of $\mathbf{H}_\mu^{-1}$ by identifying η with $(\eta_{\omega_0}, \eta_{-\omega_0})$, where, for any smooth function $\psi : \mathbf{S} \to \mathbf{R}$, $\eta_{\omega_0}(\psi) := \eta(\psi, 0)$ and $\eta_{-\omega_0}(\psi) := \eta(0, \psi)$. Defining analogously $W_{\pm\omega_0}$ for the Wiener process W in (1.9), the object of interest is then

$$\forall t > 0, \ \eta_t = \eta_0 + \int_0^t L\eta \, \mathrm{d}s + W_t. \tag{5.1}$$

5.1. **The noise W as a cylindrical Brownian Motion.** We first focus on the regularity properties of the noise W: in the stationary case ($q_t = q_t|_{t=0} = q$ for all $t > 0$) the covariance defined in (1.9) becomes, for any regular functions on $\mathbf{S} \times \mathbf{R}$, φ_1 and φ_2, $s, t > 0$:

$$\mathbf{E}\left(W_s(\varphi_1)W_t(\varphi_2)\right) = (s \wedge t) \int_{\mathbf{R}} \int_{\mathbf{S}} \partial_\theta \varphi_1 \partial_\theta \varphi_2 q \,\mathrm{d}\lambda \,\mathrm{d}\mu. \tag{5.2}$$

Consequently, it is easily seen that $(W_{+\omega_0,t}, W_{-\omega_0,t})$ is a couple of two independent Gaussian processes with covariance (where $\psi_1, \psi_2 : \mathbf{S} \to \mathbf{R}$):

$$\forall \omega \in \{+\omega_0, -\omega_0\},\ \mathbf{E}\left(W_{\omega,s}(\psi_1)W_{\omega,t}(\psi_2)\right) = \frac{1}{2}(s \wedge t) \int_{\mathbf{S}} \partial_\theta \psi_1 \partial_\theta \psi_2 q(\cdot, \omega). \tag{5.3}$$

In what follows, we will denote by $\mathbf{H}_0$ the closed subspace of $\mathbf{H}_\mu^{-1}$ consisting of elements of $\mathbf{H}_\mu^{-1}$ with zero mean-value; in particular the norm $\|\cdot\|_{\mathbf{H}_\mu}$ defined in (2.8) coincides on $\mathbf{H}_0$ with:

$$\forall h \in \mathbf{H}_0,\ \| h \|_{\mathbf{H}_\mu} = \left(2\pi \int_{\mathbf{R}} \int_{\mathbf{S}} \mathcal{H}^2\right)^{\frac{1}{2}}, \tag{5.4}$$

where we recall that $\mathcal{H}$ is the primitive of h such that $\int_{\mathbf{S}} \mathcal{H} = 0$. The purpose of this paragraph is to show that, following [9, p. 96], W has the same law as a Q-Wiener process in the Hilbert space $\mathbf{H}_0$, for an appropriate bounded symmetric operator Q on $\mathbf{H}_0$: indeed, if one denotes by X a Q-Wiener process on $\mathbf{H}_0$, with the following definition of Q

$$\forall h \in \mathbf{H}_0,\ Qh := \partial_\theta\left(q\mathcal{H}\right), \tag{5.5}$$

then one readily verifies that the Gaussian process $(W(\varphi))_\varphi$ has the same law as $(X(\varphi))_\varphi := \left(\left\langle X_t, \begin{pmatrix}\partial_\theta^2\varphi \\ 0\end{pmatrix}\right\rangle_{\mathbf{H}_\mu}, \left\langle X_t, \begin{pmatrix}0 \\ \partial_\theta^2\varphi\end{pmatrix}\right\rangle_{\mathbf{H}_\mu}\right)_\varphi$.

Note that if the weight q had not been present in (5.3), it would have been straightforward to see that W is a Q-Wiener process on $\mathbf{H}_0$, with $Q = I$. The fact that this weight is present entails some technical complications, but one has to consider the operator Q defined in (5.5) only as a perturbation of the case $Q = I$.

5.2. **Existence and uniqueness of a weak solution to the fluctuation equation.** We now turn to the existence and uniqueness of a weak solution of (5.1). We recall that any $\mathbf{H}_\mu^{-1}$-valued predictable process η_t, $t \in [0, T]$ is a *weak solution* of (5.1) if the trajectories of η are almost surely Bochner integrable and if for all $\varphi \in \mathcal{D}(L^*)$ and for all $t \in [0, T]$

$$\eta_t(\varphi) = \eta_0(\varphi) + \int_0^t \eta_s(L^*\varphi)\,\mathrm{d}s + W_t(\varphi). \tag{5.6}$$

Proposition 5.1. *Equation has a unique weak solution in* $\mathbf{H}_\mu^{-1}$, *given by the mild formulation*

$$\eta_t = T_L(t)\eta_0 + \int_0^t T_L(t-s)\,\mathrm{d}W_s, \quad t \in [0, T]. \tag{5.7}$$

To prove Proposition 5.1, one needs to define properly the stochastic convolution $W_L(t) := \int_0^t T_L(t-s)\,\mathrm{d}W_s$. In this purpose, let use prove firstly that the inverse of A is of class trace:

Lemma 5.2. *The operator* A^{-1} *is of class trace in* $\mathbf{H}_\mu^{-1}$. *Equivalently, if* $(\lambda_n^{(A)})_{n \geqslant 1}$ *is the sequence of eigenvalues of the self-adjoint operator* A, *one has*

$$\sum_{n=1}^{\infty} \frac{1}{\lambda_n^{(A)}} < \infty. \tag{5.8}$$

Proof of Lemma 5.2. Since A and $\tilde{A} = M \circ A \circ M^{-1}$ (recall (4.5)) are conjugate, it suffices to prove (5.8) when A is replaced by $\tilde{A}$. The idea of the proof is that identity (5.8) is true when $\tilde{A} = (\tilde{A}_1, \tilde{A}_2)$ is replaced by $(-\Delta, -\Delta)$ and that $\tilde{A}$ is only a relatively-bounded perturbation of this case. More precisely, the proof relies on the following MinMax Principle [13, p. 1543]:

Proposition 5.3. *Let $(F, \mathcal{D}(F))$ a self-adjoint linear operator on a separable Hilbert space $\mathcal{H}$ such that F is positive, with compact resolvent. We denote by $\mathcal{S}^n$ the family of n-dimensional subspace of $\mathcal{H}$, and for $n \geqslant 1$ we let λ_n the number defined as follows*

$$\lambda_n := \sup_{G \in \mathcal{S}^n} \inf_{u \in (G \cap \mathcal{D}(F)) \setminus \{0\}} \frac{\langle u\,,\, Fu \rangle_{\mathcal{H}}}{\langle u\,,\, u \rangle_{\mathcal{H}}}. \tag{5.9}$$

Then there exists a complete orthonormal system $(\psi_n)_{n \geqslant 1}$ such that

$$F\psi_n = \lambda_n \psi_n, \quad n \geqslant 1.$$

In other words, the sequence $(\lambda_n)_{n \geqslant 1}$ is the non-decreasing enumeration of the eigenvalues of F, each repeated a number of times equal to its multiplicity. Moreover, the sup *in* (5.9) *is attained for G equal to the span of $\{\psi_1, \ldots, \psi_n\}$.*

Let us apply Proposition 5.3 to $F = -\Delta$ with domain

$$\mathcal{D}(-\Delta) := \left\{ u;\, u \in \mathcal{C}^2(\mathbf{S}),\ \int_{\mathbf{S}} u(\theta)\, \mathrm{d}\theta = 0 \right\}, \tag{5.10}$$

in $\mathbf{H}^{-1}$ (recall the definition of $\left(\mathbf{H}^{-1}, \langle \cdot\,,\, \cdot \rangle_{-1}\right)$ in Remark 2.2) and let us denote by $\mathcal{E}_0(u, v) := \langle u\,,\, -\Delta v \rangle_{-1} = 2\pi \int_{\mathbf{S}} uv$ the Dirichlet form associated to $-\Delta$. Note that $\mathcal{E}_0$ is well defined on $\mathbf{L}^2 \supset \mathcal{D}(-\Delta)$. Then, denoting by $(\lambda_n^{(-\Delta)})_{n \geqslant 1}$ the sequence of eigenvalues associated to $-\Delta$ in $\mathbf{H}^{-1}$:

$$\lambda_n^{(-\Delta)} = \sup_{G \in \mathcal{S}^n} \inf_{u \in (G \cap \mathcal{D}(-\Delta)) \setminus \{0\}} \frac{\mathcal{E}_0(u, u)}{\langle u\,,\, u \rangle_{-1}}.$$

Since the supremum is attained for $G = \{\psi_1, \ldots, \psi_n\} \subseteq \mathbf{L}^2$, one has in fact:

$$\lambda_n^{(-\Delta)} = \sup_{G \in \mathcal{S}^n} \inf_{u \in (G \cap \mathbf{L}^2) \setminus \{0\}} \frac{\mathcal{E}_0(u, u)}{\langle u\,,\, u \rangle_{-1}}.$$

Secondly, note that one does not change the result by considering $1 - A_1$ instead of $-A_1$. Hence, if ones denotes by $\mathcal{E}_1(u, v) := \langle u\,,\, (1 - A_1)v \rangle_{-1, q_0}$ the Dirichlet form associated to $1 - A_1$, one deduces from [5, Eq.(2.47)] that $\mathcal{E}_1$ is well defined on $\mathbf{L}^2$ and that it is equivalent to $\mathcal{E}_0$: there exists a constant $C > 0$ such that

$$\forall u \in \mathbf{L}^2, \quad \frac{1}{C}\mathcal{E}_0(u, u) \leqslant \mathcal{E}_1(u, u) \leqslant C\mathcal{E}_0(u, u).$$

Then, using again Proposition 5.3,

$$\lambda_n^{(1 - A_1)} = \sup_{G \in \mathcal{S}^n} \inf_{u \in (G \cap \mathbf{L}^2) \setminus \{0\}} \frac{\mathcal{E}_1(u, u)}{\langle u\,,\, u \rangle_{-1, q_0}}.$$

Since the norms $\| \cdot \|_{-1}$ and $\| \cdot \|_{-1, q_0}$ are equivalent, one directly sees that there exist constants $c, C > 0$ such that, for all $n \geqslant 1$

$$c\lambda_n^{(-\Delta)} \leqslant \lambda_n^{(1 - A_1)} \leqslant C\lambda_n^{(-\Delta)}. \tag{5.11}$$

One can prove similar bounds for A_2 in the Hilbert space $\mathbf{H}_w^{-1}$ in the same way: first notice that any eigenvector which corresponds to a non-zero eigenvalue of A_2 is necessarily with zero mean-value, so that it suffices to work on the domain $\{v \in \mathbf{L}^2,\ \int_{\mathbf{S}} v = 0\}$. It is then easy to deduce from (4.24) that both Dirichlet forms $\mathcal{E}_0$ and $\mathcal{E}_2$ (recall definition (4.23)) are equivalent on the subspace of $\mathbf{L}^2$ with zero mean-value. In particular using Proposition 5.3, one easily obtains similar bounds as (5.11) for A_2 and (5.8) follows. □

Following the lines of [9], one can deduce that the linear operator $\int_0^t T_L(s) Q T_L(s)^* \, \mathrm{d}s$ is of class trace: indeed, it is easy to see from the definition (4.2) of the perturbation B that B satisfies the assumption (5.59) in [9, p.145], namely, B is a continuous linear operator from $\mathbf{L}^2_\mu$ into $\mathbf{H}^{-1}_\mu$, and there exists a constant $c > 0$ such that for all $h \in \mathcal{D}$, $\langle Bh\,,\, h\rangle_{\mathbf{H}_\mu} \leqslant c \| h \|^2_{\mathbf{H}_\mu}$. Since Lemma 5.2 is true, the assumptions of [9, Prop. 5.25, p.145] are satisfied, so that the operator $\int_0^t T_L(s) Q T_L(s)^* \, \mathrm{d}s$ is of class trace. Then an application of [9, Th. 5.2] shows that the stochastic convolution $W_L(\cdot)$ is well defined as a predictable process in $\mathbf{H}^{-1}_\mu$. The assumptions of [9, Th. 5.4] concerning the existence and uniqueness a weak solution of (5.1) are satisfied and Proposition 5.1 is proved.

5.3. **Linear asymptotic behavior of the fluctuation process.** We are in position to prove the main statement of Theorem 2.9, that is the asymptotic behavior (2.19) of the mild solution (5.7). We will make a constant use of the spectral decomposition of the operator L (under the hypothesis of Theorems 2.5 and 2.7). We will refer in particular to the matrix representation (2.17) of L.

Note also that the continuous linear form $\ell_{\partial_\theta q}(\cdot)$ on $\mathbf{H}^{-1}_\mu$ can be represented, by Riesz representation theorem, as a scalar product w.r.t. some vector $\zeta \in \mathbf{H}^{-1}_\mu$:

$$\ell_{\partial_\theta q}(\cdot) = \langle \zeta\,,\, \cdot \rangle_{\mathbf{H}_\mu}\,. \tag{5.12}$$

The convergence (2.19) is an easy consequence of Remark 2.8 and the following two propositions:

Proposition 5.4. *The stochastic convolution $W_L(t)$ satisfies the following linear behavior, as $t \to +\infty$: $\frac{W_L(t)}{t} \to 0$, where the convergence is in law.*

Proposition 5.5. *For every initial condition η_0, $\frac{T_L(t)\eta_0}{t}$ converges, as $t \to +\infty$, to $\ell_p(\eta_0)\partial_\theta q$.*

Let us now prove these two propositions:

Proof of Proposition 5.4. Recall that W is a Q-Wiener process in $\mathbf{H}_0$, which can be decomposed into $\mathbf{H}_0 = \mathrm{Span}(\partial_\theta q) \oplus G_{<0}$. Note also that the restriction on $\mathbf{H}_0$ of the projection P_0 defined on $\mathbf{H}^{-1}_\mu$ by (2.16) coincides with $\ell_{\partial_\theta q}(\cdot)\partial_\theta q$. With a small abuse of notations, we will use the same notation P_0 for this restriction on $\mathbf{H}_0$. Let us decompose the stochastic convolution into $W_L(t) = \int_0^t T_L(t-s) P_0 \, \mathrm{d}W_s + \int_0^t T_L(t-s) P_{<0} \, \mathrm{d}W_s$, and treat the two terms separately:

Let us consider the first term $\frac{1}{t}\int_0^t T_L(t-s) P_0 \, \mathrm{d}W_s$. One has successively, using that $T_L(u)\partial_\theta q = \partial_\theta q$ for all $u > 0$

$$\frac{1}{t}\int_0^t T_L(t-s) P_0 \, \mathrm{d}W_s = \frac{\partial_\theta q}{t}\int_0^t T_L(t-s)\ell_{\partial_\theta q} \, \mathrm{d}W_s, \tag{5.13}$$

$$= \frac{\partial_\theta q}{t}\ell_{\partial_\theta q} W_t = \frac{\partial_\theta q}{t}\langle \zeta\,,\, W_t\rangle_{\mathbf{H}_\mu}\,,\ \text{(recall (5.12))}. \tag{5.14}$$

Thanks to the Q-Wiener structure of W (see § 5.1), one has

$$\mathbf{E}\left(\left\| \frac{1}{t}\int_0^t T_L(t-s)P_0\,\mathrm{d}W_s \right\|_{\mathbf{H}_\mu}^2\right) = \frac{\|\partial_\theta q\|_{\mathbf{H}_\mu}^2}{t^2}\mathbf{E}\left(|\langle \zeta\,,\,W_t\rangle_{\mathbf{H}_\mu}|^2\right) = \frac{\|\partial_\theta q\|_{\mathbf{H}_\mu}^2}{t}\langle Q\zeta\,,\,\zeta\rangle_{\mathbf{H}_\mu}\,, \tag{5.15}$$

which converges to 0 as $t\to+\infty$.

As far as the second term $\int_0^t T_L(t-s)P_{<0}\,\mathrm{d}W_s$ is concerned, it is easy to see that it is the unique weak solution in $\mathbf{H}_0$ of

$$w_t = \int_0^t Lw_s\,\mathrm{d}s + P_{<0}\,W_t. \tag{5.16}$$

Let us decompose evolution (5.16) along this decomposition $\mathbf{H}_0 = \mathrm{Span}(\partial_\theta q)\oplus G_{<0}$: writing $w_t = P_0\,w_t + P_{<0}\,w_t := y_t + z_t$, one has:

$$\begin{cases} z_t &= \int_0^t P_0\,Ly_s\,\mathrm{d}s,\\ y_t &= \int_0^t P_{<0}\,LP_{<0}\,y_s\,\mathrm{d}s + P_{<0}\,W_t. \end{cases} \tag{5.17}$$

Since the operator $P_{<0}\,LP_{<0}$ has its spectrum in the negative part of the complex plane with a strictly positive spectral gap $\lambda_K(L)$ and generates an semigroup of operators, it is immediate to see from the covariance estimates of stochastic convolutions (see [9, Th. 5.2, p.119]) that there exist some $t_0>0$ and a constant $c>0$ such that for all $t\geqslant t_0$

$$\mathbf{E}\left(\|y_t\|_{\mathbf{H}_\mu}\right)^2 \leqslant \mathbf{E}\left(\|y_t\|_{\mathbf{H}_\mu}^2\right) \leqslant ce^{-\frac{\lambda_K(L)}{2}t} \tag{5.18}$$

Consequently, one has

$$\begin{aligned}
\mathbf{E}\left(\|z_t\|_{\mathbf{H}_\mu}\right) &\leqslant \int_0^t \mathbf{E}\left(\|P_0\,Ly_s\|_{\mathbf{H}_\mu}\right)\mathrm{d}s = \|\partial_\theta q\|_{\mathbf{H}_\mu}\int_0^t \mathbf{E}\left(|\ell_{\partial_\theta q}(Ly_s)|\right)\mathrm{d}s,\\
&= \int_0^t \mathbf{E}\left(|\langle\zeta\,,\,Ly_s\rangle_{\mathbf{H}_\mu}|\,\|\partial_\theta q\|_{\mathbf{H}_\mu}\right)\mathrm{d}s,\ \text{(recall (5.12))}\\
&\leqslant \|\partial_\theta q\|_{\mathbf{H}_\mu}\|L^*\zeta\|_{\mathbf{H}_\mu}\int_0^t \mathbf{E}\left(\|y_s\|_{\mathbf{H}_\mu}\right)\mathrm{d}s,\\
&= \|\partial_\theta q\|_{\mathbf{H}_\mu}\|L^*\zeta\|_{\mathbf{H}_\mu}\left(\int_0^{t_0}\mathbf{E}\left(\|y_s\|_{\mathbf{H}_\mu}\right)\mathrm{d}s + \int_{t_0}^t \mathbf{E}\left(\|y_s\|_{\mathbf{H}_\mu}\right)\mathrm{d}s\right).
\end{aligned} \tag{5.19}$$

In particular, it is immediate from (5.18) and (5.19) to see that the following convergence holds:

$$\frac{\mathbf{E}\left(\|z_t\|_{\mathbf{H}_\mu}\right)}{t}\to_{t\to+\infty} 0. \tag{5.20}$$

Putting together (5.18) and (5.20), Proposition 5.4 is proved. □

Proof of Proposition 5.5. Let us fix an initial condition $\eta_0\in\mathbf{H}_\mu^{-1}$. Then $X(t) := T_L(t)\eta_0$ is the unique solution in $\mathbf{H}_\mu^{-1}$ of

$$X(t) = \eta_0 + \int_0^t LX_s\,\mathrm{d}s. \tag{5.21}$$

Decompose $X(t)$ along the direct sum $G_0\oplus G_{<0}$, that is $X(t) = \alpha(t)\partial_\theta q + \beta(t)p + Y(t)$, with $Y(t)\in G_{<0}$. Then, projecting on $\partial_\theta q$, p and $G_{<0}$ respectively (see (2.17)), one obtains that

(5.21) is equivalent to

$$\forall t > 0, \quad \begin{cases} \alpha(t) &=& \ell_{\partial_\theta q}(\eta_0) + \int_0^t \left(\beta(s) + \ell_{\partial_\theta q}(LP_{<0} Y(s))\right) \mathrm{d}s, \\ \beta(t) &=& \ell_p(\eta_0), \\ Y(t) &=& P_{<0}\eta_0 + \int_0^t P_{<0} L P_{<0} Y(s) \,\mathrm{d}s. \end{cases} \tag{5.22}$$

Then, since $T_{P_{<0} L P_{<0}}(t)$ is a semigroup of contraction whose infinitesimal generator has a strictly positive spectral gap $\lambda_K(L)$, there exists in particular a constant $c > 0$ such that $Y(t) = T_{P_{<0} L P_{<0}}(t) P_{<0}\eta_0$ and $\| Y(t) \|_{\mathbf{H}_\mu} \leqslant c e^{-\frac{\lambda_K(L)}{2} t}$ (in particular, $\frac{1}{t} \| Y(t) \|_{\mathbf{H}_\mu} \to_{t \to +\infty} 0$). Then, using again (5.12),

$$\frac{\alpha(t)}{t} = \frac{\ell_{\partial_\theta q}(\eta_0)}{t} + \ell_p(\eta_0) + \frac{1}{t} \int_0^t \ell_{\partial_\theta q}(L P_{<0} Y(s)) \,\mathrm{d}s, \tag{5.23}$$

$$= \frac{\ell_{\partial_\theta q}(\eta_0)}{t} + \ell_p(\eta_0) + \frac{1}{t} \int_0^t \langle P_{<0}^* L^* \zeta \,,\, Y(s) \rangle_{\mathbf{H}_\mu} \,\mathrm{d}s. \tag{5.24}$$

Using the previous exponential bound for $Y(s)$, it is easy to see that $\frac{\alpha(t)}{t}$ converges to $\ell_p(\eta_0)$ as $t \to +\infty$. The result of Proposition 5.5 follows. □

Appendix A. Gelfand-triple construction

The construction of the weighted-Sobolev spaces defined in § 2.2 and used throughout the paper is based on the usual notion of Gelfand-triple (or rigged-Hilbert spaces) that we precise here. We refer to [5, § 2.2] or [7, p.81] for precise definitions.

Namely, one can understand the closure of $\left\{h \in \mathcal{C}^1(\mathbf{S});\ \int_{\mathbf{S}} h = 0\right\}$ w.r.t. the norm $\|h\|_{-1,k}$ defined in (2.7) as the dual space V' of the space V closure of $\left\{h \in \mathcal{C}^1(\mathbf{S});\ \int_{\mathbf{S}} h = 0\right\}$ with respect to the norm

$$\|h\|_V := \left(\int_{\mathbf{S}} h'(\theta)^2 k(\theta) \,\mathrm{d}\theta\right)^{\frac{1}{2}}.$$

The pivot space is the usual $\mathbf{L}^2(\lambda)$, endowed with the Hilbert norm

$$\|h\|_{\mathbf{L}^2} := \left(\int_{\mathbf{S}} h(\theta)^2 \,\mathrm{d}\theta\right)^{\frac{1}{2}}.$$

In particular, one easily sees that the inclusion $V \subseteq \mathbf{L}^2(\lambda)$ is dense. Consequently, one can define $T : \mathbf{L}^2(\lambda) \to V'$ by setting $Th(v) = \int_{\mathbf{S}} h(\theta) v(\theta) \,\mathrm{d}\theta$. One can prove that T continuously injects $\mathbf{L}^2(\lambda)$ into V' and that $T(\mathbf{L}^2(\lambda))$ is dense into V' so that one can identify $h \in \mathbf{L}^2(\lambda)$ with $Th \in V'$. Then for $h \in \mathbf{L}^2(\lambda)$,

$$\|h\|_{V'} = \|Th\|_{V'} = \sup_{v \in V} \frac{\int_{\mathbf{S}} \mathcal{H} h'}{\|v\|_V} = \sqrt{\int_{\mathbf{S}} \frac{\mathcal{H}^2}{k}}, \tag{A.1}$$

where we used in (A.1) Cauchy-Schwarz inequality for the lower bound and chose $v' := \frac{\mathcal{H}}{k}$ for the upper bound. This enables to identify $\mathbf{H}_k^{-1}$ with V'.

Acknowledgements

This is a part of my PhD thesis. I would like to thank my PhD supervisors Giambattista Giacomin and Lorenzo Zambotti for introducing this subject, for their useful advice, and for their encouragement. I am also indebted to Christophe Poquet for useful discussions.

References

[1] J. A. Acebrón, L. L. Bonilla, C. J. Pérez Vicente, F. Ritort, and R. Spigler. The Kuramoto model: A simple paradigm for synchronization phenomena. *Rev. Mod. Phys.*, 77(1):137–185, Apr 2005.

[2] D. G. Aronson. Addendum: "Non-negative solutions of linear parabolic equations" (Ann. Scuola Norm. Sup. Pisa (3) **22** (1968), 607–694). *Ann. Scuola Norm. Sup. Pisa (3)*, 25:221–228, 1971.

[3] P. B. Bailey, W. N. Everitt, and A. Zettl. Regular and singular Sturm-Liouville problems with coupled boundary conditions. *Proc. Roy. Soc. Edinburgh Sect. A*, 126(3):505–514, 1996.

[4] N. J. Balmforth and R. Sassi. A shocking display of synchrony. *Physica D: Nonlinear Phenomena*, 143(1-4):21–55, 2000.

[5] L. Bertini, G. Giacomin, and K. Pakdaman. Dynamical aspects of mean field plane rotators and the Kuramoto model. *J. Statist. Phys.*, 138:270–290, 2010.

[6] P. Billingsley. *Probability and measure*. Wiley Series in Probability and Mathematical Statistics. John Wiley & Sons Inc., New York, third edition, 1995.

[7] H. Brezis. *Analyse fonctionnelle*. Collection Mathématiques Appliquées pour la Maîtrise. [Collection of Applied Mathematics for the Master's Degree]. Masson, Paris, 1983. Théorie et applications. [Theory and applications].

[8] E. A. Coddington and N. Levinson. *Theory of ordinary differential equations*. McGraw-Hill Book Company, Inc., New York-Toronto-London, 1955.

[9] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*, volume 44 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1992.

[10] P. Dai Pra and F. den Hollander. McKean-Vlasov limit for interacting random processes in random media. *J. Statist. Phys.*, 84(3-4):735–772, 1996.

[11] F. den Hollander. *Large deviations*, volume 14 of *Fields Institute Monographs*. American Mathematical Society, Providence, RI, 2000.

[12] N. Dunford and J. T. Schwartz. *Linear operators. Part II*. Wiley Classics Library. John Wiley & Sons Inc., New York, 1988. Spectral theory. Selfadjoint operators in Hilbert space, With the assistance of William G. Bade and Robert G. Bartle, Reprint of the 1963 original, A Wiley-Interscience Publication.

[13] N. Dunford and J. T. Schwartz. *Linear operators. Part III*. Wiley Classics Library. John Wiley & Sons Inc., New York, 1988. Spectral operators, With the assistance of William G. Bade and Robert G. Bartle, Reprint of the 1971 original, A Wiley-Interscience Publication.

[14] M. S. P. Eastham. Results and problems in the spectral theory of periodic differential equations. In *Spectral theory and differential equations (Proc. Sympos., Dundee, 1974; dedicated to Konrad Jörgens)*, pages 126–135. Lecture Notes in Math., Vol. 448. Springer, Berlin, 1975.

[15] A. Friedman. *Partial differential equations of parabolic type*. Prentice-Hall Inc., Englewood Cliffs, N.J., 1964.

[16] J. Gärtner. On the McKean-Vlasov limit for interacting diffusions. *Math. Nachr.*, 137:197–248, 1988.

[17] G. Giacomin, E. Luçon, and C. Poquet. Coherence stability and effect of random natural frequencies in population of coupled oscillators, 2011. `arxiv.org/abs/1111.3581`.

[18] G. Giacomin, K. Pakdaman, and X. Pellegrin. Global attractor and asymptotic dynamics in the Kuramoto model for coupled noisy phase oscillators, 2011. `arxiv.org/abs/1107.4501`.

[19] G. Giacomin, K. Pakdaman, X. Pellegrin, and C. Poquet. Transitions in active rotator systems: invariant hyperbolic manifold approach, 2011. `arxiv.org/abs/1106.0758`.

[20] E. Hille and R. S. Phillips. *Functional analysis and semi-groups*. American Mathematical Society Colloquium Publications, vol. 31. American Mathematical Society, Providence, R. I., 1957. rev. ed.

[21] B. Jourdain and F. Malrieu. Propagation of chaos and Poincaré inequalities for a system of particles interacting through their CDF. *Ann. Appl. Probab.*, 18(5):1706–1736, 2008.

[22] B. Jourdain and S. Méléard. Propagation of chaos and fluctuations for a moderate model with smooth initial data. *Ann. Inst. H. Poincaré Probab. Statist.*, 34(6):727–766, 1998.

[23] T. Kato. *Perturbation theory for linear operators*. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the 1980 edition.

[24] Y. Kuramoto. Self-entrainment of a population of coupled non-linear oscillators. In *International Symposium on Mathematical Problems in Theoretical Physics (Kyoto Univ., Kyoto, 1975)*, pages 420–422. Lecture Notes in Phys., 39. Springer, Berlin, 1975.

[25] Y. Kuramoto. *Chemical oscillations, waves, and turbulence*, volume 19 of *Springer Series in Synergetics*. Springer-Verlag, Berlin, 1984.

[26] E. Luçon. Quenched limits and fluctuations of the empirical measure for plane rotators in random media. *Electronic Journal of Probability*, 16:792–829, 2011.

[27] A. Lunardi. *Analytic semigroups and optimal regularity in parabolic problems*. Progress in Nonlinear Differential Equations and their Applications, 16. Birkhäuser Verlag, Basel, 1995.

[28] A. S. Markus. *Introduction to the spectral theory of polynomial operator pencils*, volume 71 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1988. Translated from the Russian by H. H. McFaden, Translation edited by Ben Silver, With an appendix by M. V. Keldysh.

[29] H. P. McKean, Jr. Propagation of chaos for a class of non-linear parabolic equations. In *Stochastic Differential Equations (Lecture Series in Differential Equations, Session 7, Catholic Univ., 1967)*, pages 41–57. Air Force Office Sci. Res., Arlington, Va., 1967.

[30] M. Möller and A. Zettl. Symmetric differential operators and their Friedrichs extension. *J. Differential Equations*, 115(1):50–69, 1995.

[31] M. A. Naĭmark. *Linear differential operators. Part II: Linear differential operators in Hilbert space*. With additional material by the author, and a supplement by V. È. Ljance. Translated from the Russian by E. R. Dawson. English translation edited by W. N. Everitt. Frederick Ungar Publishing Co., New York, 1968.

[32] K. Oelschläger. A martingale approach to the law of large numbers for weakly interacting stochastic processes. *Ann. Probab.*, 12(2):458–479, 1984.

[33] A. Pazy. *Semigroups of linear operators and applications to partial differential equations*, volume 44 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1983.

[34] P. A. Pearce. Mean-field bounds on the magnetization for ferromagnetic spin models. *J. Statist. Phys.*, 25(2):309–320, 1981.

[35] H. Risken. *The Fokker-Planck equation*, volume 18 of *Springer Series in Synergetics*. Springer-Verlag, Berlin, second edition, 1989. Methods of solution and applications.

[36] H. Sakaguchi. Cooperative phenomena in coupled oscillator systems under external fields. *Progr. Theoret. Phys.*, 79(1):39–46, 1988.

[37] S. Shinomoto and Y. Kuramoto. Phase transitions in active rotator systems. *Progress of Theoretical Physics*, 75(5):1105–1110, 1986.

[38] R. E. Showalter. *Monotone operators in Banach space and nonlinear partial differential equations*, volume 49 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997.

[39] S. H. Strogatz. From Kuramoto to Crawford: exploring the onset of synchronization in populations of coupled oscillators. *Phys. D*, 143(1-4):1–20, 2000. Bifurcations, patterns and symmetry.

[40] S. H. Strogatz and R. E. Mirollo. Stability of incoherence in a population of coupled oscillators. *J. Statist. Phys.*, 63(3-4):613–635, 1991.